



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 9/10, 15/54, 15/63, 15/70, 15/82, 5/10, A01H 5/00, C12P 21/02, 19/04, 19/18</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/47727</b> <b>(43) International Publication Date:</b> 17 August 2000 (17.08.00)
<b>(21) International Application Number:</b> PCT/EP00/00954 <b>(22) International Filing Date:</b> 7 February 2000 (07.02.00) <b>(30) Priority Data:</b> 199 05 069.4      8 February 1999 (08.02.99)      DE <b>(71) Applicants (for all designated States except US):</b> PLANT-TEC BIOTECHNOLOGIE GMBH FORSCHUNG & ENTWICKLUNG [DE/DE]; Hermannswerder 14, D-14473 Potsdam (DE). MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN e.V. [DE/DE]; Berlin (DE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> KOSSMANN, Jens [DE/DE]; Düppelstr. 29, D-14163 Berlin (DE). WELSH, Thomas [DE/DE]; Schönburgstrasse 20, D-12103 Berlin (DE). QUANZ, Martin [DE/DE]; Oppelner Strasse 34, D-10997 Berlin (DE). KNUTH, Karola [DE/DE]; Schöneberger Str. 6, D-12103 Berlin (DE). <b>(74) Agent:</b> VOSSIUS & PARTNER; Siebertstrasse 4, D-81765 München (DE).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>

**(54) Title:** NUCLEIC ACID MOLECULES ENCODING ALTERNANSUCRASE**(57) Abstract**

Nucleic acid molecules encoding an alternansucrase are provided. Moreover, vectors, host cells and plant cells transformed by the herein-described nucleic acid molecules and plants containing them are provided. Furthermore, methods are described for preparing transgenic plants which synthesize the carbohydrate alternan, because of the insertion of nucleic acid molecules encoding an alternansucrase. Moreover, methods for preparing alternan and products resulting from them are provided.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

### **Nucleic Acid Molecules Encoding Alternansucrase**

The present invention relates to nucleic acid molecules encoding an alternansucrase. Moreover, this invention relates to vectors, host cells and plant cells transformed with the herein-described nucleic acid molecules, and plants containing said cells. Moreover, methods for preparing transgenic plants which due to the insertion of DNA molecules encoding an alternansucrase, synthesize the carbohydrate alternan, are described. Furthermore, methods for preparing alternan are described.

Prior art documents, the disclosure content of which is included into the present application by reference thereto, are cited hereinafter.

Alternan is a polysaccharide composed of glucose units. The glucose units are linked to each other via  $\alpha$ -1,3- and  $\alpha$ -1,6-glycosidic bonds, and said two types of bonds predominantly appear alternatingly. However, alternan is not a linear polysaccharide, but may contain branches (Seymour et al., Carbohydrate Research 74, (1979), 41-62). Because of its physico-chemical properties, the possibilities of application of alternan both in the pharmaceutical industry, for instance as a carrier of pharmaceutically active ingredients and as an additive in the textile, cosmetics and food industry have been discussed (Lopez-Munguia et al., Enzyme Microb. Technol. 15, (1993), 77-85; Leathers et al., Journal of Industrial Microbiology & Biotechnology 18, (1997), 278-283). Moreover, it can be used as a substitute for gum arabic (Coté, Carbohydrate Polymers 19, (1992), 249-252).

Industry has a high interest in biotechnological methods for preparing oligosaccharides and polysaccharides, and in particular alternan which is hardly or not at all accessible to classical organic synthesis. Compared to the classical approach of organic synthesis chemistry, biotechnological processes offer

advantages. For instance, enzymatically catalyzed reactions as a rule show much higher specificities (regio specificity, stereo specificity) and higher reaction speeds, proceed under milder reaction conditions and lead to higher yields. These factors are of outstanding importance in the preparation of new oligosaccharides and polysaccharides.

Alternan is prepared enzymatically with the use of enzymes possessing the biological activity of alternansucrases. Alternansucrases belong to the group of glucosyltransferases, which, starting from saccharose, are able to catalyze the formation of alternan and fructose. So far, alternansucrases have only been found in the bacterium *Streptococcus mutans* (Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063); Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353)) and in specific strains of the gram positive bacterium *Leuconostoc mesenteroides* where they are, as a rule, present together with other polysaccharide-forming enzymes, such as for instance dextran-forming dextransucrases, or together with polysaccharide-degrading enzymes, such as alternanases. Hence, the naturally occurring strains also produce dextran in addition to alternan.

So far, alternan has been prepared in a cell-free system using partially purified proteins or by fermentation using alternansucrase-producing strains of *Leuconostoc mesenteroides*.

Various purification methods for the purification of alternansucrases have been described (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85; Lopez-Munguia et al., Annals New York Academy of Sciences 613 (1990), 717-722; Côté and Robyt, Carbohydrate Research 101 (1982), 57-74). These methods are complex and relatively costly, and, as a rule, lead to low protein yields (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283). None of these methods allows highly pure alternansucrase protein to be produced, and therefore sequencing of the protein and the isolation of the corresponding DNA sequences have not been successful so far. If the alternansucrase protein purified according to these methods is used for *in vitro* preparation of alternan, then the dextransucrase protein residues contained in the alternansucrase preparation produce dextran impurities in the alternan produced. The separation of alternan and dextran is relatively time-consuming and costly (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283). Another disadvantage of the

dextranucrase protein impurities contained in the enzyme preparation of alternansucrase protein is the fact that a part of the saccharose substrate is converted into dextran and not into alternan, which results in a reduction of the alternan yield.

The fermentative preparation by means of *Leuconostoc* also leads to the formation of product mixtures of alternan and dextran. In order to increase the amount of alternansucrase from *Leuconostoc* strains, mutants have been isolated, such as the mutant NRRL B-21138, which secrete the alternansucrase and lead to a higher proportion of the amount of alternansucrase formed relative to dextranucrase. However, if such mutants are fermented with sucrose, the alternan obtained continues to show dextran impurities (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283).

As can be seen from the prior art discussed above, it has not been possible to provide highly purified alternansucrase protein so far.

Hence, the present invention addresses the problem of providing means and methods allowing alternan to be prepared in a time-saving and inexpensive manner.

This problem is solved by the provision of the embodiments characterized in the patent claims.

Consequently, the present invention relates to a nucleic acid molecule encoding a protein possessing the biological activity of an alternansucrase selected from the group consisting of

- (a) nucleic acid molecules encoding at least the mature form of a protein which comprises the amino acid sequence indicated in Seq. ID No. 2 or the amino acid sequence encoded by the cDNA contained in plasmid DSM 12666;
- (b) nucleic acid molecules comprising the nucleotide sequence indicated in Seq. ID No. 1 or the nucleotide sequence of the cDNA contained in plasmid DSM 12666 or a corresponding ribonucleotide sequence;

- (c) nucleic acid molecules encoding a protein, the amino acid sequence of which has a homology of at least 40% to the amino acid sequence indicated in Seq. ID No. 2;
- (d) nucleic acid molecules, one strand of which hybridizes with the nucleic acid molecules as defined in (a) or (b);
- (e) nucleic acid molecules comprising a nucleotide sequence encoding a biologically active fragment of the protein which is encoded by any one of the nucleic acid molecules as defined in (a), (b), (c) or (d); and
- (f) nucleic acid molecules, the nucleotide sequence of which deviates because of the degeneration of the genetic code from the sequence of the nucleic acid molecules as defined in (a), (b), (c), (d) or (e).

Consequently, the present invention relates to nucleic acid molecules encoding proteins possessing the biological activity of an alternansucrase, said molecules preferably encoding proteins comprising the amino acid sequence indicated in Seq. ID No. 2.

An enzyme possessing the enzymatic or biological activity of an alternansucrase (E.C. 2.4.1.140) is understood to mean an enzyme which is able to catalyze the conversion of saccharose into alternan and fructose. This conversion may occur both in the presence and absence of external acceptors (for instance maltose, isomaltose, isomaltotriose etc.). In the absence of external acceptors, alternansucrases starting from saccharose catalyze the release of fructose and high molecular alternan, a polysaccharide composed of glucose units, the backbone of which consists of glucose units predominantly connected to each other alternately by  $\alpha$ -1,3- and  $\alpha$ -1,6-glycosidic bonds. Concerning the percentage of  $\alpha$ -1,3- and  $\alpha$ -1,6-linked glucose units the literature displays different values. According to Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063), alternan consists of 76 mol%  $\alpha$ -1,3-linked glucose and 24 mol%  $\alpha$ -1,6-linked glucose. Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353) describe alternan as a polyglucan containing 49.1 mol%  $\alpha$ -1,6-linked glucose and 33.9 mol%  $\alpha$ -1,3-linked glucose with 13.6 mol% terminal glucose and 3.3 mol%  $\alpha$ -1,3,6-branched glucose. In the presence of external acceptors, such as maltose, isomaltose, isomaltotriose and

methyl- $\alpha$ -D-glucan, alternansucrase can catalyze the synthesis of  $\alpha$ -D-glucan chains, in which the glucose residues are predominantly alternatingly connected by  $\alpha$ -1,6- and  $\alpha$ -1,3-glycosidic bonds, and the synthesis of fructose at these polysaccharide acceptors. Depending on the acceptor used, the products formed have different structures. The enzymatic activity of an alternansucrase can for instance be detected as described by Lopez-Munguia (Annals New York Academy of Sciences 613 (1990), 717-722) or as described in the examples of the present application.

The invention in particular relates to nucleic acid molecules containing the nucleotide sequence indicated under Seq. ID No. 1 or a part thereof, and preferably to molecules, which comprise the coding region indicated in Seq. ID No. 1 or corresponding ribonucleotide sequences.

Moreover, the present invention relates to nucleic acid molecules which encode an alternansucrase and the one strand of which hybridizes with one of the above-described molecules.

The present invention also relates to nucleic acid molecules which encode a protein, which has a homology, that is to say an identity of at least 40%, preferably at least 60%, preferably at least 70%, especially preferably at least 80% and in particular at least 90% to the entire amino acid sequence indicated in Seq. ID No. 2, the protein possessing the biological activity of an alternansucrase.

The present invention also relates to nucleic acid molecules, which encode an alternansucrase and the sequence of which deviates on account of the degeneration of the genetic code from the nucleotide sequences of the above-described nucleic acid molecules.

The invention also relates to nucleic acid molecules possessing a sequence which is complementary to the whole or a part of the above-mentioned sequences.

The nucleic acid sequence indicated in Seq. ID No. 1 for instance encodes an extracellular alternansucrase. Secretion is ensured by a signal sequence which comprises the first approximately 39 N-terminal amino acid groups of the Seq. ID No. 2. In certain circumstances it may be desirable for only the mature protein to be expressed without naturally occurring signal sequences and/or together with other

signal sequences. Hence, the above-described nucleic acid molecules encode at least the mature form of a protein possessing the biological activity of an alternansucrase.

Within the present invention the term "hybridization" means hybridization under conventional hybridization conditions, preferably under stringent conditions, as for instance described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2<sup>nd</sup> edition (1989) Cold Spring Harbour Laboratory Press, Cold Spring Harbour, NY. Within an especially preferred meaning the term "hybridization" means that hybridization occurs under the following conditions:

Hybridization buffer:	2 x SSC; 10 x Denhardt solution (Fikoll 400 + PEG + BSA; ratio 1:1:1); 0.1% SDS; 5 mM EDTA; 50 mM Na <sub>2</sub> HPO <sub>4</sub> ; 250 µg/ml of herring sperm DNA; 50 µg/ml of tRNA; or 0.25 M of sodium phosphate buffer, pH 7.2; 1 mM EDTA 7% SDS
Hybridization temperature T	= 60°C
Washing buffer:	2 x SSC; 0.1% SDS
Washing temperature T	= 60°C.

Nucleic acid molecules which hybridize with the nucleic acid molecules of the invention can, in principle, encode alternansucrases from any organism expressing such proteins.

Nucleic acid molecules which hybridize with the molecules of the invention can for instance be isolated from genomic libraries of microorganisms. Alternatively, they can be prepared by genetic engineering or chemical synthesis.

Such nucleic acid molecules may be identified and isolated with the use of the molecules of the invention or parts of these molecules or reverse complements of these molecules, for instance by hybridization according to standard methods (see for instance Sambrook et al., 1989, Molecular Cloning. A Laboratory Manual, 2<sup>nd</sup> edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

Nucleic acid molecules possessing the same or substantially the same nucleotide sequence as indicated in Seq. ID No. 1 or parts thereof can, for instance, be used



as hybridization probes. The fragments used as hybridization probes can also be synthetic fragments which are prepared by usual synthesis techniques, and the sequence of which substantially coincides with that of an inventive nucleic acid molecule.

The molecules hybridizing with the nucleic acid molecules of the invention also comprise fragments, derivatives and allelic variants of the above-described nucleic acid molecules encoding an alternansucrase of the invention. Herein, fragments are understood to mean parts of the nucleic acid molecules which are long enough to encode one of the described proteins, preferably showing the biological activity of an alternansucrase. In this connection, the term derivative means that the sequences of these molecules also differ from the sequences of the above-described nucleic acid molecules in one or more positions and show a high degree of homology to these sequences. In this context, homology means a sequence identity of at least 40%, in particular an identity of at least 60%, preferably more than 80% and particularly preferably more than 90%. Deviations from the above-described nucleic acid molecules may have been produced by deletion, substitution, insertion and/or recombination.

Preferably, the degree of homology is determined by comparing the respective sequence with the nucleotide sequence of the coding region of SEQ ID No.1. When the sequences which are compared do not have the same length, the degree of homology preferably refers to the percentage of nucleotide residues in the shorter sequence which are identical to nucleotide residues in the longer sequence. The degree of homology can be determined conventionally using known computer programs such as the ClustalW program (Thompson et al., Nucleic Acids Research 22 (1994), 4673-4680) distributed by Julie Thompson (Thompson@EMBL-Heidelberg.DE) and Toby Gibson (Gibson@EMBL-Heidelberg.DE) at the European Molecular Biology Laboratory, Meyerhofstrasse 1, D 69117 Heidelberg, Germany. ClustalW can also be downloaded from several websites including IGBMC (Institut de Génétique et de Biologie Moléculaire et Cellulaire, B.P.163, 67404 Illkirch Cedex, France; <ftp://ftp-igbmc.u-strasbg.fr/pub/>) and EBI (<ftp://ftp.ebi.ac.uk/pub/software/>) and all sites with mirrors to the EBI (European

Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SD, UK).

When using ClustalW program version 1.8 to determine whether a particular sequence is, for instance, 90% identical to a reference sequence according to the present invention, the settings are set in the following way for DNA sequence alignments:

KTUPLE=2, TOPDIAGS=4, PAIRGAP=5, DNAMATRIX:IUB, GAOPEN=10, GAPEXT=5, MAXDIV=40, TRANSITIONS: unweighted.

For protein sequence alignments using ClustalW program version 1.8 the settings are the following: KTUPLE=1, TOPDIAG=5, WINDOW=5, PAIRGAP=3, GAOPEN=10, GAPEXTEND=0.05, GAPDIST=8, MAXDIV=40, MATRIX=GONNET, ENDGAPS(OFF), NOPGAP, NOHGAP.

Furthermore, homology means preferably that the encoded protein displays a sequence identity of at least 40%, more preferably of at least 60%, even more preferably of at least 80%, in particular of at least 90% and particularly preferred of at least 95% to the amino acid sequence depicted under SEQ ID NO: 2.

Homology, moreover, means that there is a functional and/or structural equivalence between the corresponding nucleic acid molecules or proteins encoded thereby. Nucleic acid molecules which are homologous to the above-described molecules and represent derivatives of these molecules are, as a rule, variations of these molecules which represent modifications having the same biological function. They may be either naturally occurring variations, for instance sequences from other microorganisms, or mutations, and said mutations may have formed naturally or may have been produced by deliberate mutagenesis. Furthermore, the variations may be synthetically produced sequences. The allelic variants may be naturally occurring variants or synthetically produced variants or variants produced by recombinant DNA techniques.

In a further preferred embodiment the term "derivative" encompasses a nucleic acid molecule coding for a protein which comprises at least one, more preferably at least three, even more preferably at least five, in particular at least ten and particularly preferred at least twenty of the peptide motifs selected from the group consisting of

- a) MKQQE (SEQ ID NO: 22),
- b) KKVPPV (SEQ ID NO: 23),

- c) KDDEN (SEQ ID NO: 24),
- d) IDGNL (SEQ ID NO: 25),
- e) YVADS (SEQ ID NO: 26),
- f) HLRKN (SEQ ID NO: 27),
- g) NENTP (SEQ ID NO: 28),
- h) NVDGY (SEQ ID NO: 29),
- i) NPDLK (SEQ ID NO: 30),
- j) SNDSG (SEQ ID NO: 31),
- k) NTFVK (SEQ ID NO: 32),
- l) ISGYL (SEQ ID NO: 33),
- m) SNAAL (SEQ ID NO: 34),
- n) RQYTD (SEQ ID NO: 35),
- o) QLYRA (SEQ ID NO: 36),
- p) DDKAP (SEQ ID NO: 37),
- q) TRQYT (SEQ ID NO: 38),
- r) ITFAG (SEQ ID NO: 39),
- s) NQYKG (SEQ ID NO: 40),
- t) LFLNA (SEQ ID NO: 41),
- u) QVSDT (SEQ ID NO: 42),
- v) LITLN (SEQ ID NO: 43),
- w) GRYVH (SEQ ID NO: 44),
- x) TAPYG (SEQ ID NO: 45),
- y) VVDYQ (SEQ ID NO: 46),
- z) LSGQE (SEQ ID NO: 47).

The proteins encoded by the different variants of the nucleic acid molecules of the invention possess certain characteristics they have in common. These include for instance enzymatic activity, molecular weight, immunological reactivity, conformation, etc., and physical properties, such as for instance the migration behavior in gel electrophoreses, chromatographic behavior, sedimentation coefficients, solubility, spectroscopic properties, stability, pH optimum, temperature optimum etc.

Alternansucrase (E.C. 2.4.1.140) is an enzyme belonging to the group of glucosyltransferases. So far, alternansucrase activity has not been found in plants, but only in the bacterium *Streptococcus mutans* (Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063); Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353)) and in specific strains of the bacterium *Leuconostoc mesenteroides*, for instance in NRRL B-1355, NRRL B-1498 and NRRL B-1501. As a rule, these strains contain different glucosyltransferases and secrete dextransucrases apart from alternansucrases if they are allowed to grow on saccharose-containing media. As a rule, these two sucrases possess a high binding affinity to the polysaccharides synthesized by them (Lopez-Munguia et al., Annals New York Academy of Sciences 613 (1990), 717-722) with the result that these polysaccharides must be separated from the protein in the purification of the enzymes from *Leuconostoc mesenteroides* strains grown on saccharose-containing medium (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85; Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283).

In the absence of external acceptors, alternansucrases, starting from saccharose, catalyze the release of fructose and high molecular alternan, a polysaccharide which is composed of glucose units, and the backbone of which consists of glucose units predominantly linked to each other alternatingly by  $\alpha$ -1,3- and  $\alpha$ -1,6-glycosidic bonds and which according to light scattering measurement data should have a molecular weight of  $>10^7$  (Coté, Carbohydrate Polymer 19 (1992), 249-252). To date there has been no report of alternan possessing a terminal fructose residue. Nevertheless, the existence of a terminal fructose unit in alternan can not be completely excluded. Lopez-Munguia et al. (Enzyme Microb. Technol. 15 (1993) 77-85) describe that alternan is resistant to degradation by dextranases. However, it can be degraded by so-called alternanases, whereby ring-shaped oligomers of alternan of different polymerization degree can be produced (Biely et al., Eur. J. Biochem. 226 (1994), 633-639). Ultrasonic treatment of high molecular alternan allows the molecular weight of alternan to be reduced to  $<10^6$  (Coté, Carbohydrate Polymers 19 (1992), 249-252). If aqueous solutions of this ultrasonically treated alternan are prepared, then these solutions show rheological properties comparable to those of aqueous solutions of gum arabic. So-called "limit alternan" having a molecular weight of about 3500 can be produced by enzymatic degradation using

isomaltodextranase from *Arthrobacter globiformis* (NRRL B-4425) (Coté, Carbohydrate Polymers 19 (1992), 249-252).

In the presence of external acceptors, such as for instance maltose, isomaltose, isomaltotriose and methyl- $\alpha$ -D-glucan, alternansucrase catalyzes at said saccharide acceptors the synthesis of  $\alpha$ -D-glucan chains, in which the glucose moieties are predominantly alternatingly linked by  $\alpha$ -1,6- and  $\alpha$ -1,3 glycosidic bonds, and the synthesis of fructose. Depending on the acceptor used, the resulting products have different structures and a molecular weight which is lower than that of high molecular alternan and a polymerization degree of  $<15$ . Because of the polymerization degree, these products are often also referred to as oligoalternans (Pelenc et al., Sciences Des Aliments 11 (1991), 465-476). However, within the framework of the present invention these low molecular products which can be prepared in the presence of external acceptors are also to be referred to as alternan.

In the preparation of oligoalternans by means of partially purified alternansucrase protein, maltose is an acceptor (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85) producing high oligoalternan yields. Panose (degree of polymerization (d.p.) of 3) is the first acceptor product which is formed starting from maltose through the formation of an  $\alpha$ -1,6-glycosidic bond.

In contrast thereto, isomaltose is a less effective acceptor which leads to lower yields of oligoalternan (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

Alternansucrase is relatively stable and has a half life period of 2 days in 50 mM of acetate buffer, pH 5.4 at 40°C (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85). The enzyme shows maximum activity at a temperature of 40°C and a pH value of 5.6 (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

In the absence of the substrate saccharose, alternansucrase catalyzes disproportionation reactions leading to a (partial) rearrangement of alternan. In particular when partially purified alternansucrase preparations containing dextransucrase contaminations were used to prepare oligoalternans, high disproportionation rates were found which lead to a complete rearrangement of oligoalternan (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

For the molecular weight of alternansucrase according to SDS PAGE determination, different numerical values can be found: 135 kDa, 145 kDa, 173 kDa and 196 kDa, respectively (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283; Kim & Robyt, Enzyme Microb. Technol. 16 (1994), 659-664; Zhanley & Smith, Applied and Environmental Microbiology 61(3) (1995), 1120-1123).

The enzymatic activity of an alternansucrase can be shown for instance as described in Lopez-Munguia et al. (Annals New York Academy of Sciences 613 (1990), 717-722) or as described in the examples of the present application.

One activity unit (1u) can be defined as the amount of enzyme leading to the release of 1  $\mu$ mol of fructose within one minute.

The nucleic acid molecules of the invention can be DNA molecules, in particular genomic molecules. Moreover, the nucleic acid molecules of the invention may be RNA molecules. The nucleic acid molecules of the invention can be obtained for instance from natural sources or may be produced synthetically or by recombinant techniques.

The nucleic acid molecules of the invention allow host cells to be prepared which produce recombinant alternansucrase protein of high purity and/or in sufficient quantities, and genetically engineered plants possessing an activity of these enzymes leading to the formation of alternan *in planta*. Within the framework of the present invention the term "high purity" means that the protein according to the invention displays a degree of purity of at least 80%, preferably of at least 90%, even more preferably of at least 95%. Moreover, means and methods are provided which may be used for preparing alternan using host cells and/or for preparing recombinant alternansucrase protein. Consequently, the provision of the nucleic acid molecules of the invention permits the preparation of alternan of high purity by methods which are relatively inexpensive and consume relatively little time.

In a preferred embodiment, the nucleic molecules of the invention are derived from microorganisms, preferably from bacteria, more preferably from gram-positive bacteria and in particular preferably from bacteria belonging to the genus

*Leuconostoc*. Nucleic acid molecules from bacteria belonging to the species *Leuconostoc mesenteroides* are particularly preferred.

The invention also relates to oligonucleotides specifically hybridizing to a nucleic acid molecule of the invention. Such oligonucleotides have a length of preferably at least 10, in particular at least 15, and particularly preferably of at least 50 nucleotides. They are characterized in that they specifically hybridize to the nucleic acid molecules of the invention, that is to say that they do not or only to a very minor extent hybridize to nucleic acid sequences encoding other proteins, in particular other glucosyltransferases. The oligonucleotides of the invention can be used for instance as primers for amplification techniques such as the PCR reaction or as a hybridization probe to isolate related genes.

Moreover, the invention relates to vectors, in particular plasmids, cosmids, viruses, bacteriophages and other vectors commonly used in gene technology, which contain the above-described nucleic acid molecules of the invention. In a preferred embodiment of the invention, the vectors of the invention lend themselves to the transformation of fungal cells or cells of microorganisms. Preferably, such vectors are suitable to transform plant cells. Particularly preferably, such vectors permit the integration of the nucleic acid molecules of the invention, possibly together with flanking regulatory regions, into the genome of the plant cell. Examples thereof are binary vectors which can be used in the *Agrobacteria*-mediated gene transfer, and some are already commercially available.

In another preferred embodiment, the nucleic acid molecules contained in the vectors are connected to regulatory elements ensuring the transcription and synthesis of a translatable RNA in prokaryotic or eukaryotic cells.

The expression of the nucleic acid molecules of the invention in prokaryotic or eukaryotic cells, for instance in *Escherichia coli*, is interesting because it permits a more precise characterization of the enzymatic activities of the enzymes encoded by these molecules. Moreover, it is possible to express these enzymes in such prokaryotic or eukaryotic cells which are free from interfering enzymes, such as dextranases or other polysaccharide-forming or polysaccharide-degrading

enzymes. In addition, it is possible to insert different mutations into the nucleic acid molecules by methods usual in molecular biology (see for instance Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2<sup>nd</sup> edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY), leading to the synthesis of proteins possibly having modified biological properties. On the one hand it is possible in this connection to produce deletion mutants in which nucleic acid molecules are produced by progressive deletions from the 5' or 3' end of the coding DNA sequence, and said nucleic acid molecules lead to the synthesis of correspondingly shortened proteins. Such deletions at the 5' end of the nucleotide sequence for instance allow amino acid sequences to be identified which are responsible for the secretion of the enzyme in microorganisms (transit peptides).

This permits the deliberate preparation of enzymes which are no longer secreted by the removal of the corresponding sequences, but remain within the cell of the corresponding host organism or are localized in other compartments, for instance in the plastids, mitochondria, vacuole, on account of the addition of other signal sequences.

On the other hand, the introduction of point mutations is also conceivable at positions at which a modification of the amino acid sequence for instance influences the enzyme activity or the control of the enzyme. In this manner, it is for instance possible to produce mutants which possess a modified stereo and regio selectivity or a modified  $K_m$  value or which are no longer subject to the control mechanisms normally existing in the cell and realized via an allosteric control or covalent modification.

Moreover, mutants possessing a modified substrate or product specificity can be prepared. Furthermore, it is possible to prepare mutants having a modified activity-temperature-profile.

Furthermore, in the case of expression in plants, the insertion of mutations into the nucleic acid molecules of the invention allows the gene expression rate and/or the activity of the proteins encoded by the nucleic acid molecules of the invention to be increased.

For genetic engineering in prokaryotic cells, the nucleic acid molecules of the invention or parts of these molecules can be introduced into plasmids which permit mutagenesis or sequence modification by recombination of DNA sequences.



Standard methods (see Sambrook et al., 1989, Molecular Cloning: A laboratory manual, 2<sup>nd</sup> edition, Cold Spring Harbor Laboratory Press, NY, USA) allow base exchanges to be performed or natural or synthetic sequences to be added. DNA fragments can be connected to each other by applying adapters and linkers to the fragments. Moreover, engineering measures which provide suitable restriction sites or remove surplus DNA or restriction sites can be used. In those cases, in which insertions, deletions or substitutions are possible, *in vitro* mutagenesis, "primer repair", restriction or ligation can be used. In general, a sequence analysis, restriction analysis and other methods of biochemistry and molecular biology are carried out as analysis methods.

Moreover, the invention relates to plasmid pAlsu-pSK (see Fig. 2 and Example 2) which was deposited at Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ), Braunschweig, under the accession No. DSM 12666 on February 4, 1999, and to the nucleic acid molecules contained in the insert of plasmid DSM 12666 and encoding a protein possessing the enzymatic activity of an alternansucrase. Moreover, the present invention also relates to nucleic acid molecules which hybridize to the insertion of plasmid DSM 12666. Also, the present invention relates to nucleic acid molecules the nucleotide sequence of which deviates from that of the nucleic acid molecules of the plasmid DSM 12666 insert, because of the degeneration of the genetic code. Furthermore, the present invention relates to nucleic acid molecules which have a homology, that is to say a sequence identity of at least 40%, preferably of at least 60%, more preferably of at least 80%, even more preferably of at least 90%, and most preferably of at least 95% to the sequence of the insertion of plasmid DSM 12666.

Another embodiment of the invention relates to host cells, in particular prokaryotic or eukaryotic cells transformed with an above-described nucleic acid molecule of the invention or with a vector of the invention, and to cells descended from such transformed cells and containing a nucleic acid molecule or vector of the invention. According to another preferred embodiment, the host cells are cells of microorganisms. In the context of the present invention, the term "microorganism" comprises bacteria and all protists (e.g. fungi, in particular yeasts, algae) as defined

Schlegel's "Allgemeine Mikrobiologie" (Georg Thieme Verlag, 1985, 1-2). A preferred embodiment of the invention relates to cells of algae and host cells belonging to the genera *Aspergillus*, *Bacillus*, *Saccharomyces* or *Pichia* (Rodriguez, Journal of Biotechnology 33 (1994), 135-146, Romanos, Vaccine, Vol. 9 (1991), 901 et seq.). A particularly preferred embodiment of the invention relates to *E. coli* cells. Alternansucrase is especially preferably secreted by the host cell. The preparation of such host cells for the production of recombinant alternansucrase can be carried out by methods known to a man skilled in the art.

In a preferred embodiment of the invention, the host cells of the invention show no interfering enzymatic activities, such as those of polysaccharide-forming and/or polysaccharide-degrading enzymes.

An overview of different expression systems is for instance contained in Methods in Enzymology 153 (1987), 385-516, in Bitter et al. (Methods in Enzymology 153 (1987), 516-544) and in Sawers et al. (Applied Microbiology and Biotechnology 46 (1996), 1-9), Billman-Jacobe (Current Opinion in Biotechnology 7 (1996), 500-4), Hockney (Trends in Biotechnology 12 (1994), 456-463), Griffiths et al, Methods in Molecular Biology 75 (1997), 427-440). An overview of yeast expression systems is for instance given by Hensing et al. (Antonie van Leeuwenhoek 67 (1995), 261-279), Bussineau et al. (Developments in Biological Standardization 83 (1994), 13-19), Gellissen et al. (Antonie van Leeuwenhoek 62 (1992), 79-93, Fleer (Current Opinion in Biotechnology 3 (1992), 486-496), Vedvick (Current Opinion in Biotechnology 2 (1991), 742-745) and Buckholz (Bio/Technology 9 (1991), 1067-1072).

Expression vectors have been widely described in the literature. As a rule, they contain not only a selection marker gene and a replication-origin ensuring replication in the host selected, but also a bacterial or viral promoter, and in most cases a termination signal for transcription. Between the promoter and the termination signal there is at least one restriction site or a polylinker which enables the insertion of a coding DNA sequence. The DNA sequence naturally controlling the transcription of the corresponding gene can be used as the promoter sequence, if it is active in the selected host organism. However, this sequence can also be exchanged for other promoter sequences. It is possible to use promoters producing a constitutive expression of the gene and inducible promoters which permit a deliberate control of the expression of the postconnected gene. Bacterial and viral

promoter sequences possessing these properties are described in detail in the literature. Regulatory sequences for the expression in microorganisms (for instance *E. coli*, *S. cerevisiae*) are sufficiently described in the literature. Promoters permitting a particularly high expression of the postconnected gene are for instance the T7 promoter (Studier et al., Methods in Enzymology 185 (1990), 60-89), lacUV5, trp, trp-lacUV5 (DeBoer et al., in Rodriguez and Chamberlin (Eds), Promoters, Structure and Function; Praeger, New York, (1982), 462-481; DeBoer et al., Proc. Natl. Acad. Sci. USA (1983), 21-25), lp1, rac (Boros et al., Gene 42 (1986), 97-100). As a rule, the protein amounts are highest from the middle up to about the end of the logarithmic phase of the growth cycle of the microorganisms. Therefore, inducible promoters are preferably used for the synthesis of proteins. These promoters often lead to higher protein yields than do constitutive promoters. The use of highly constitutive promoters leads to the continuous transcription and translation of a cloned gene and thus often has the result that energy is lost for other essential cells functions with the effect that cell growth is slowed down (Bernard R. Glick/Jack J. Pasternak, Molekulare Biotechnologie (1995). Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin, Oxford, p. 342). Therefore, in order to obtain an optimum amount of protein, a two-stage process is often used. First, the host cells are cultured under optimum conditions up to a relatively high cell density. In the second step, transcription is then induced depending on the type of promoter used. In this connection, a tac promoter is particularly suitable which can be induced by lactose or IPTG (=isopropyl- $\beta$ -D-thiogalactopyranoside) (deBoer et al., Proc. Natl. Acad. Sci. USA 80 (1983), 21-25). Termination signals for transcription are also described in the literature.

The transformation of the host cell with DNA encoding an alternansucrase can, as a rule, be carried out by standard methods, as for instance described in Sambrook et al., (Molecular Cloning: A Laboratory Course Manual, 2<sup>nd</sup> edition (1989) Cold Spring Harbor Press, New York; Methods in Yeast Genetics, A Laboratory Course Manual, Cold Spring Harbor Laboratory Press, 1990). The host cell is cultured in nutrient media meeting the requirements of the particular host cell used, in particular in respect of the pH value, temperature, salt concentration, aeration, antibiotics, vitamins, trace elements etc.

Moreover, the invention relates to proteins and biologically active fragments thereof, which are encoded by the nucleic acid molecules of the invention and to methods for their preparation, wherein a host cell according to the invention is cultured under conditions permitting the synthesis of the protein, and the protein is subsequently isolated from the cultured cells and/or the culture medium.

According to a preferred embodiment of the invention, the alternansucrase is a recombinantly produced protein. In the context of the present invention, this is a protein prepared by inserting a DNA sequence encoding the protein into a host cell and expressing it therein. The protein can then be isolated from the host cell and/or the culture medium.

The nucleic acid molecules of the invention now allow host cells to be prepared which produce recombinant alternansucrase protein of high purity and/or in sufficient amounts. Within the framework of the present invention the term "high purity" means that the protein according to the invention displays a degree of purity of at least 80%, preferably of at least 90%, even more preferably of at least 95%. The time-consuming and costly methods already mentioned above, whereby alternansucrase protein which to date can only be obtained from particular *Leuconostoc* strains can be purified from other components such as for instance dextransucrases, polysaccharides, are dispensed with, because alternansucrase can be produced in host cells not possessing any adverse polysaccharide-synthesizing activities. Moreover, host cells and vectors can also be used, which allow the alternansucrase protein to be produced in the absence of saccharose, with the result that an additional separation of the alternansucrase protein from polysaccharides is no longer necessary. Moreover, the selection of suitable host cells and vectors allows alternansucrase protein to be provided in sufficient amounts, which has not been possible with the systems so far described.

Alternansucrase produced by the host cells can be purified by conventional purification methods, such as precipitation, ion exchange chromatography, affinity-chromatography, gel filtration, HPLC Reverse Phase Chromatography etc.

The modification of the nucleic acid molecules of the invention encoding an alternansucrase and expressed in the host cells, allows to produce a polypeptide in the host cell which is easier to isolate from the culture medium because of particular

properties. Thus, the protein to be expressed can be expressed as a fusion protein with an additional polypeptide sequence, the specific binding properties of which permit the isolation of the fusion protein by affinity chromatography (e.g. Hopp et al., Bio/Technology 6 (1988), 1204-1210; Sassenfeld, Trends Biotechnol. 8 (1990), 88-93).

Another embodiment of the invention relates to proteins possessing the enzymatic activity of an alternansucrase, in particular that from microorganisms, preferably Gram-positive microorganisms, particularly microorganisms of the genus *Leuconostoc*, and particularly preferably that from *Leuconostoc mesenteroides*. The molecular weight of the protein indicated in Seq. ID No. 2, as determined by calculation, is 228.96 kDa. The invention also relates to alternansucrases which possess a molecular weight of  $229 \text{ kDa} \pm 120 \text{ kDa}$ , preferably  $229 \text{ kDa} \pm 50 \text{ kDa}$ , and particularly preferably  $230 \text{ kDa} \pm 25 \text{ kDa}$ . The molecular weight of the mature protein, as determined by calculation, is 224.77 kDa.

The provision of the nucleic acid molecules of the invention, for the first time, makes it possible to prepare alternansucrase-expressing plant cells by means of genetic engineering, which was not possible so far, because classical culturing methods do not allow bacterial and fungal genes to be expressed in plants.

The invention, therefore, also relates to transgenic plant cells transformed by a nucleic acid molecule of the invention or a vector of the invention or descended from such cells, the nucleic acid molecule which encodes the protein that has the biological activity of an alternansucrase being under the control of regulatory elements permitting the transcription of a translatable mRNA in plant cells.

The introduction of the activity of the proteins of the invention, for instance by expression of corresponding nucleic acid molecules, opens the possibility of producing alternan in plant cells correspondingly modified by genetic engineering. Hence, the expression of the nucleic acid molecules of the invention in plant cells is possible, allowing an additional, corresponding alternansucrase activity not present in the wild type to be introduced. Moreover, it is possible to modify the nucleic acid molecules of the invention according to methods known to a skilled person, in order to obtain alternansucrases of the invention which for instance possess modified

temperature dependencies or substrate or product specificities. Such methods have already been described in more detail in a different context above.

A plurality of techniques is available by which DNA can be inserted into a plant host cell. These techniques include the transformation of plant cells by T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as a transforming agent, the fusion of protoplasts, injection, electroporation of DNA, insertion of DNA by the biolistic approach and other possibilities.

The use of the Agrobacteria-mediated transformation of plant cells has been extensively investigated and sufficiently described in EP 120 516; Hoekema, In: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblasterdam (1985), Chapter V; Fraley et al, Crit. Rev. Plant Sci. 4 (1993), 1-46 and An et al., EMBO J. 4 (1985), 277-287. Regarding the transformation of potatoes see for instance Rocha-Sosa et al. (EMBO J. 8 (1989), 29-33).

The transformation of monocotyledonous plants by means of Agrobacterium-based vectors has also been described (Chan et al., Plant Mol. Biol. 22 (1993), 491-506; Hiei et al., Plant J. 6 (1994) 271-282; Deng et al, Science in China 33 (1990), 28-34; Wilmerk et al, Plant Cell Reports 11 (1992), 76-80; May et al., Bio/Technology 13 (1995), 486-492; Conner and Dormisse, Int. J. Plant Sci. 153 (1992), 550-555; Ritchie et al. Transgenic Res. 2 (1993), 252-265). An alternative system for transforming monocotyledonous plants is the transformation by the biolistic approach (Wan and Lemaux, Plant Physiol. 104 (1994), 37-48; Vasil et al., Bio/Technology 11 (1993), 1553-1558; Ritala et al., Plant Mol. Biol. 24 (1994) 317-325; Spencer et al., Theor. Appl. Genet. 79 (1990), 625-631), protoplast transformation, electroporation of partially permeabilized cells, insertion of DNA via glass fibers. The transformation of maize in particular has been repeatedly described in the literature (see for instance WO 95/06128, EP 0 513 849, EP 0 465 875, EP 29 24 35; Fromm et al, Biotechnology 8, (1990), 833-844; Gordon-Kamm et al., Plant Cell 2, (1990), 603-618; Koziel et al., Biotechnology 11 (1993), 194-200; Moroc et al., Theor. Appl. Genet. 80, (1990), 721-726).

The successful transformation of other types of cereals has also been described for instance of barley (Wan and Lemaux, supra; Ritala et al., supra, Krens et al., Nature 296 (1982), 72-74) and wheat (Nehra et al., Plant J. 5 (1994), 285-297). Generally, any promoter active in plant cells is suitable to express the nucleic acid

molecules in plant cells. The promoter can be so chosen that the expression in the plants of the invention occurs constitutively or only in a particular tissue, at a particular time of plant development or at a time determined by external influences. The promoter may be homologous or heterologous to the plant.

Suitable promoters are for instance the promoter of 35S RNA of the Cauliflower Mosaic Virus (see for instance US-A-5,352,605) and the ubiquitin-promoter (see for instance US-A-5,614,399) which lend themselves to constitutive expression, the patatin gene promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29) which lends itself to a tuber-specific expression in potatoes or a promoter ensuring expression in photosynthetically active tissues only, for instance the ST-LS1 promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO, J. 8 (1989) 2445-2451), the Ca/b-promoter (see for instance US-A-5,656,496, US-A-5,639,952, Bansal et al., Proc. Natl. Acad. Sci. USA 89 (1992), 3654-3658) and the Rubisco SSU promoter (see for instance US-A-5,034,322; US-A-4,962,028) or the glutelin promoter from wheat which lends itself to endosperm-specific expression (HMW promoter) (Anderson, Theoretical and Applied Genetics 96, (1998), 568-576, Thomas, Plant Cell 2 (12), (1990), 1171-1180), the glutelin promoter from rice (Takaiwa, Plant Mol. Biol. 30(6) (1996), 1207-1221, Yoshihara, FEBS Lett. 383 (1996), 213-218, Yoshihara, Plant and Cell Physiology 37 (1996), 107-111), the shrunken promoter from maize (Maas, EMBO J. 8 (11) (1990), 3447-3452, Werr, Mol. Gen. Genet. 202(3) (1986), 471-475, Werr, Mol. Gen. Genet. 212(2), (1988), 342-350), the USP promoter, the phaseolin promoter (Sengupta-Gopalan, Proc. Natl. Acad. Sci. USA 82 (1985), 3320-3324, Bustos, Plant Cell 1 (9) (1989), 839-853) or promoters of zein genes from maize (Pedersen et al., Cell 29 (1982), 1015-1026; Quatroccio et al., Plant Mol. Biol. 15 (1990), 81-93). However, promoters which are only activated at a point in time determined by external influences can also be used (see for instance WO 93/07279). In this connection, promoters of heat shock proteins which permit simple induction may be of particular interest. Moreover, seed-specific promoters such as the USP promoter from *Vicia faba* which ensures a seed-specific expression in *Vicia faba* and other plants may be used (Fiedler et al., Plant Mol. Biol. 22 (1993), 669-679; Bäumlein et al., Mol. Gen. Genet. 225 (1991), 459-467). Moreover, fruit-specific promoters, such as described in WO 91/01373 may be used too.

Moreover, a termination sequence may be present, which serves to terminate transcription correctly and to add a poly-A-tail to the transcript, which is believed to have a function in the stabilization of the transcripts. Such elements are described in the literature (see for instance Gielen et al., EMBO J. 8 (1989), 23-29) and can be replaced at will.

Such cells can be distinguished from naturally occurring plant cells *inter alia* by the fact that they contain a nucleic acid molecule of the invention which does not naturally occur in these cells. Moreover, such transgenic plant cells of the invention can be distinguished from naturally occurring plant cells in that they contain at least one copy of the nucleic acid molecule of the invention stably integrated in their genome.

Moreover, the plant cells of the invention can preferably be distinguished from naturally occurring plant cells by at least one of the following features: If the inserted nucleic acid molecule of the invention is heterologous to the plant cell, then the transgenic plant cells are found to have transcripts of the inserted nucleic acid molecules of the invention. The latter can be detected for instance by Northern blot analysis. The plants cells of the invention preferably contain a protein encoded by an inserted nucleic acid molecule of the invention. This can be shown for instance by immunological methods, in particular by Western blot analysis.

Transgenic plant cells can be regenerated to whole plants according to methods known to a person skilled in the art.

The present invention also relates to the plants obtainable by regeneration of the transgenic plant cells of the invention. Furthermore, it relates to plants containing the above-described transgenic plant cells.

In most plants, the photoassimilates in the form of sugars formed during photosynthesis within a plant, i.e. mainly in the form of saccharose, are transported to the corresponding target organs. As saccharose is the substrate of the polymerization reaction of alternansucrase, all plants, both monocotyledonous and dicotyledonous can, in principle, be modified by the nucleic acid molecule of the invention in respect of alternansucrase expression.



The expression in plants of the nucleic acid molecules of the invention encoding a protein having the enzymatic activity of an alternansucrase can, for instance, be used to achieve a modification of the viscosity of the extracts possibly obtained from the plants, said modification being achieved by the synthesis of alternan. In this connection, for instance tomatoes are of interest. The expression of an alternansucrase in a tomato fruit leads to the synthesis of alternan and results in a modification of the viscosity of extracts obtained from these fruits for instance for the production of tomato puree or tomato ketchup.

The expression of the nucleic acid molecules of the invention is in particular advantageous in those organs of the plant which show a higher saccharose content or store saccharose. Such organs are for instance the beet of sugar beet or the cane of sugar cane. As these plants normally do not store any appreciable amounts of starch, the alternans synthesized by the alternansucrase from these plants could be isolated in the pure form.

The site where the biosynthesis of the saccharose in the plant cell occurs is the cytosol. The storage site, however, is the vacuole. During its transport into the storage tissue of the sugar beet or the potato or during its transport into the endosperm of seeds, the saccharose must pass the apoplast. Hence, all three compartments, i.e. the cytosol, the vacuole, the apoplast, lend themselves to the expression of the nucleic acid molecules for the synthesis of alternan. In addition, the plastids also lend themselves thereto, as could for instance be shown by the expression of bacterial fructosyl transferases in amyloplasts. Said fructosyl transferases which likewise require saccharose as a substrate, were able to mediate the formation of "amylofructan" in amyloplasts (Smeekens, Trends in Plant Science, Vol. 2, No. 8 (1997), 286-288).

In the case of starch-producing plants, such as potatoes and maize, where the starch biosynthesis and starch storage normally take place in the amyloplasts, an expression of the alternansucrase in apoplasts, in the cytosol or in the vacuole would lead to an additional synthesis of oligosaccharides and/or polysaccharides in these compartments, which can mean an overall increase in the yield.

As in the case of potatoes the starch synthesized in the amyloplasts can be separated from the alternan synthesized in the apoplast, in the cytosol or in the vacuole, the very same plant can be used to recover starch and alternan.

Moreover, transgenic potato and maize plants are known, the starch synthesis of which in the tubers and grains, respectively, is completely inhibited due to the inhibition of ADP-glucose-pyrophosphorylase by an antisense construct. In the case of potatoes, soluble sugars, in particular saccharose and glucose, accumulate instead, for instance in the tubers (Müller-Röber et al., EMBO J. 11 (1992), 1229-1238). Alternan can be prepared in the cytosol, the vacuole or apoplast of these plants by the expression of an alternansucrase which uses saccharose as a substrate.

Therefore in another embodiment of the invention the plant cells of the invention are further characterized by a reduced ADP glucose pyrophosphorylase (AGPase) activity compared to corresponding cells from wild-type plants.

DNA molecules encoding AGPase are well known to the person skilled in the art and described for example in Müller-Röber et al. (Mol. Gen. Genet. 224 (1) (1990), 136-146). By using DNA molecules encoding an AGPase it is possible to produce plants by means of recombinant DNA techniques (for example by an antisense, a ribozyme or a cosuppression approach) showing a reduced AGPase activity. Furthermore AGPase mutants, for example from maize (brittle-2 and shrunken-2), with reduced AGPase activity are known to the person skilled in the art.

The term "reduced" means preferably a reduction of AGPase activity of at least 10%, more preferably of at least 50% and even more preferably of at least 80% in comparison to corresponding wild-type cells.

The activity of an AGPase can be determined according to Müller-Röber et al. (Mol. Gen. Genet. 224 (1) (1990), 136-146) or to methods known to a person skilled in the art.

The reaction which is catalyzed by alternansucrase is distinguished by the fact that a glucose moiety is transferred directly from saccharose to an existing carbohydrate acceptor. By contrast, in the case of plants, the biosynthesis of linear glucans from saccharose, proceeds in such a way that the saccharose is first separated into glucose and fructose, which are then each converted into activated intermediate ADP-glucose. The glucose moiety is transferred by the enzyme starch-synthase from the ADP glucose to an already existing glucan, whereby ADP is released. The

conversion of saccharose into two ADP glucose molecules requires several energy consuming reactions. Therefore, the energy consumption of the reaction catalyzed by alternansucrase is substantially lower than the energy consumption in the synthesis of polysaccharides from saccharose in plant cells, which can lead to an increased yield of synthesized oligo and/or polysaccharides in plants containing the nucleic acid molecules of the invention.

In the expression of the nucleic acid molecules in plants there exists in principle the possibility that the synthesized protein can be localized in any compartment of the plant cell (e.g. in the cytosol, plastids, vacuole, mitochondria) or the plant (e.g. in the apoplast). In order to achieve the localization in a particular compartment, the coding region must, where necessary, be linked to DNA sequences ensuring localization in the corresponding compartment. The signal sequences used must each be arranged in the same reading frame as the DNA sequence encoding the enzyme.

In order to ensure the location in the plastids it is conceivable to use one of the following transit peptides: of the plastidic Ferredoxin: NADP<sup>+</sup> oxidoreductase (FNR) of spinach which is enclosed in Jansen et al. (Current Genetics 13 (1988), 517-522). In particular, the sequence ranging from the nucleotides -171 to 165 of the cDNA Sequence disclosed therein can be used, which comprises the 5'non-translated region as well as the sequence encoding the transit peptide. Another example is the transit peptide of the waxy protein of maize including the first 34 amino acid residues of the mature waxy protein (Klösgen et al., Mol. Gen. Genet. 217 (1989), 155-161). It is also possible to use this transit peptide without the first 34 amino acids of the mature protein. Furthermore, the signal peptides of the ribulose biphosphate carboxylase small subunit (Wolter et al., Proc. Natl. Acad. Sci. USA 85 (1988), 846-850; Nawrath et al., Proc. Natl. Acad. Sci. USA 91 (1994), 12760-12764), of the NADP malat dehydrogenase (Gallardo et al., Planta 197 (1995), 324-332), of the glutathione reductase (Creissen et al., Plant J. 8 (1995), 167-175) or of the R1 protein Lorberth et al. (Nature Biotechnology 16, (1998), 473-477) can be used.

In order to ensure the location in the vacuole it is conceivable to use one of the following transit peptides: the N-terminal sequence (146 amino acids) of the patatin

protein (Sonnewald et al., Plant J. 1 (1991), 95-106) or the signal sequences described by Matsuoka und Neuhaus, Journal of Experimental Botany 50 (1999), 165-174; Chrispeels und Raikhel, Cell 68 (1992), 613-616; Matsuoka und Nakamura, Proc. Natl. Acad. Sci. USA 88 (1991), 834-838; Bednarek und Raikhel, Plant Cell 3 (1991), 1195-1206; Nakamura und Matsuoka, Plant Phys. 101 (1993), 1-5 .

In order to ensure the location in the mitochondria it is for example conceivable to use the transit peptide described by Braun et al.(EMBO J. 11, (1992), 3219-3227).

In order to ensure the location in the apoplast it is conceivable to use one of the following transit peptides: signal sequence of the proteinase inhibitor II-gene (Keil et al., Nucleic Acid Res. 14 (1986), 5641-5650; von Schaewen et al., EMBO J. 9 (1990), 30-33), of the levansucrase gene from *Erwinia amylovora* (Geier and Geider, Phys. Mol. Plant Pathol. 42 (1993), 387-404), of a fragment of the patatin gene B33 from *Solanum tuberosum*, which encodes the first 33 amino acids (Rosahl et al., Mol Gen. Genet. 203 (1986), 214-220) or of the one described by Oshima et al. (Nucleic Acid Res. 18 (1990), 181).

The nucleic acid sequence indicated in Seq. ID No. 1 encodes an extracellular alternansucrase. Secretion is ensured by a signal sequence comprising the first approximately 39 N-terminal amino acid residues of the Seq. ID No. 2.

The transgenic plants may, in principle, be plants of any plant species, that is to say they may be monocotyledonous and dicotyledonous plants. Preferably, the plants are useful plants cultivated by man for nutrition or for technical, in particular industrial, purposes. They are preferably starch-storing plants, for instance cereal species (rye, barley, oat, wheat, millet, sago etc.), rice, pea, marrow pea, cassava and potato; tomato, rape, soybean, hemp, flax, sunflower, cow pea or arrowroot, fiber-forming plants (e.g. flax, hemp, cotton), oil-storing plants (e.g. rape, sunflower, soybean) and protein-storing plants (e.g. legumes, cereals, soybeans). The invention also relates to fruit trees and palms. Moreover, the invention relates to forage plants (e.g. forage and pasture grasses, such as alfalfa, clover, ryegrass) and vegetable plants (e.g. tomato, lettuce, chicory) and ornamental plants (e.g. tulips, hyacinths). Sugar-storing and/or starch-storing plants are preferred. Sugar

cane and sugar beet, and potato plants, maize, rice, wheat and tomato plants are particularly preferred.

A further subject of the invention is a method for the production of transgenic plant cells and transgenic plants which in comparison to non-transformed wildtype cells / non-transformed wildtype plants synthesize alternan. In this method the expression and/or the activity of proteins encoded by the nucleic acid molecules of the invention is increased in comparison to corresponding wild-type cells / wildtype plants which do not show any alternansucrase expression and/or activity. In particular, such a method comprises the expression of a nucleic acid molecule according to the invention in plant cells. The nucleic acid molecule according to the invention is preferably linked to a promoter ensuring expression in plant cells. In a particularly preferred embodiment the method comprises the introduction of a nucleic acid molecule according to the invention into a plant cell and regeneration of a plant from this cell.

Such an increase in expression may, e.g., be detected by Northern blot analysis. The increase in activity may be detected by testing protein extracts for their alternansucrase activity derived from plant cells. The enzymatic activity of an alternansucrase can be measured, for instance, as described in Lopez-Munguia et al. (Annals New York Academy of Sciences 613, (1990), 717-722) or as described in the examples of the present application.

The invention also relates to propagation material of the plants of the invention. The term "propagation material" comprises those components of the plant which are suitable to produce offspring vegetatively or generatively. Suitable means for vegetative propagation are for instance cuttings, callus cultures, rhizomes or tubers. Other propagation material includes for instance fruits, seeds, seedlings, protoplasts, cell cultures etc. The preferred propagation materials are tubers and seeds. The invention also relates to harvestable parts of the plants of the invention such as, for instance, fruits, seeds, tubers or rootstocks.

Another embodiment of the invention relates to methods for preparing alternan which comprise the step of extracting and isolating alternan from a plant of the invention.

The extraction and isolation of alternan from a plant of the invention may be carried out by standard methods, such as precipitation, extraction and chromatographic methods.

Moreover, the present invention relates to alternan obtainable from a plant of the invention or from propagation material of the invention.

Moreover, the present invention relates to a method for preparing alternan and/or fructose, wherein a host cell of the invention secretes an alternansucrase into a saccharose-containing culture medium and alternan and/or fructose is/are isolated from the culture medium.

A preferred embodiment of the method of the invention uses an alternansucrase recombinantly produced and secreted by the host cell into the culture medium, thus avoiding the necessity of breaking up the cells and purifying the protein further, because the secreted protein can be obtained from the supernatant. The residual components of the culture medium can be removed by methods usual in processing technology, such as dialysis, reverse osmosis, chromatographic methods, etc. The same applies to the concentration of the protein secreted into the culture medium. The secretion of proteins by microorganisms is normally mediated by N-terminal signal peptides (signal sequence, leader-peptide, transit peptide). Proteins possessing this signal sequence are able to penetrate the cell membrane of the microorganism. A secretion of proteins can be achieved by adding the DNA sequence encoding this signal peptide to the corresponding region encoding the alternansucrase.

The natural signal peptide of the expressed alternansucrase is preferred, that of the alternansucrase from *Leuconostoc mesenteroides* NRRL B 1355 (see the first approximately 25 to 45 N-terminal amino acid residues of Seq. ID No. 2) is particularly preferred.

The signal peptide of  $\alpha$ -CGTase from *Klebsiella oxytoca* M5A1 (Fiedler et al., J. Mol. Biol. 256 (1996), 279-291) or a signal peptide as encoded by the nucleotides 11529-11618 of the sequence available under the GenBank accession number X86014 is most preferred.

The preparation of alternan and/or fructose requires neither activated glucose derivatives nor co-factors, as are necessary in most synthesis reactions for polysaccharides occurring within the cells. Hence, alternansucrase-secreting microorganisms can be cultured in saccharose-containing medium, the secreted alternansucrase leading to a synthesis of alternan and fructose in the culture medium.

Contrary to host cells from *Leuconostoc mesenteroides*, which secrete alternansucrase by nature, the host cells used according to the invention have the advantage that they do not secrete proteins possessing adverse polysaccharide-synthesizing side reactions, such as dextransucrase, with the result that outside the host cell, apart from alternan, no other polysaccharides can be formed which, as a rule, can be separated from alternan only by costly and time-consuming procedures. Moreover, the host cells according to a preferred embodiment of the invention do not have any adverse polysaccharide-degrading side activities, which would otherwise lead to losses in the yield of the alternan produced.

The method of the invention yields fructose apart from alternan. Fructose can be used for the inexpensive isolation of so-called "high-fructose-containing syrups" (HFCS). Conventional methods for preparing fructose on the one hand provide for the enzymatic break down of saccharose by means of an invertase or for the break down of starch into glucose units, mostly brought about by acid hydrolysis, and for subsequent enzymatic conversion of the glucose into fructose by glucose isomerases. However, both methods lead to mixtures of glucose and fructose. The two components must subsequently be separated from each other by chromatographic methods.

The separation of the two reaction products of the method of the invention, or the separation of the reaction products from the substrate saccharose can be achieved for example with the use of membranes permitting the penetration of fructose, but not the penetration of saccharose and/or alternans. If continuous removal of

fructose via such a membrane is provided for, a more or less complete conversion of saccharose occurs.

The isolation of alternan and fructose can be carried out by standard methods or can be carried out as for instance described in the working examples.

According to one embodiment of the method, the host cells originate from microorganisms, preferably from *Escherichia coli*.

In another embodiment, the method of the invention works with fungal host cells, in particular cells of yeasts, such as *Saccharomyces cerevisiae*. Yeast cells producing alternan in saccharose-containing medium because of the enzymatic activity of an alternansucrase, cannot be readily used, as yeasts secrete an invertase which breaks down the extracellular saccharose. The yeasts take up the resulting hexoses via a hexose transporter. However, one yeast strain has been described (Riesmeier et al. EMBO J. 11 (1992), 4705-4713) which carries a defective *suc2* gene, and therefore cannot secrete invertase. Moreover, these yeast cells do not contain a transportation system able to import saccharose into the cells. If such a strain is so modified by means of the nucleic acid molecules of the invention that it secretes an alternansucrase into the culture medium, then fructose and alternan will be synthesized in saccharose-containing medium. The resulting fructose can subsequently be taken up by the yeast cells.

In another preferred embodiment of this method the host cell of the invention is present in an immobilized form.

As a rule, host cells are immobilized by inclusion of the cells in a suitable material, such as alginate, polyacrylamide, gelatin, cellulose or chitosan. However, adsorption or covalent binding of the cells to a carrier material is also possible (Brodellius and Mosbach, Methods in Enzymology Vol. 135 (1987), 222-230) An advantage of the immobilization of cells is that it allows substantially higher cell densities to be achieved than does culturing in liquid culture. This results in a higher productivity. Moreover, the costs for agitation and aeration of the culture decrease as do the costs for measures to maintain sterility. Another important aspect is the



possibility of a continuous alternan production with the result that unproductive phases regularly occurring in fermentation processes can be avoided or at least greatly reduced.

Another embodiment of the invention relates to a method for preparing alternan and/or fructose, wherein

- a) a saccharose-containing solution is contacted with a protein of the invention under conditions permitting the conversion of saccharose into alternan and/or fructose; and
- b) alternan and/or fructose is/are isolated from the solution.

In this embodiment, the invention thus relates to a method for preparing alternan and/or fructose *in vitro* by means of a cell-free enzyme preparation. In this case, microorganisms which for instance secrete alternansucrase are cultured up to the stationary phase in a saccharose-free medium permitting the formation of alternansucrase protein. After removing the cells from the culture medium by centrifugation, the secreted enzyme can be recovered from the supernatant. The enzyme can subsequently be added to saccharose-containing solutions in order to synthesize alternan and/or fructose. Compared to the above-described synthesis of alternan in a system not freed from cells, this method offers the advantage that the reaction conditions can be controlled better and the reaction products are substantially purer and easier to purify. The purification of the protein can be carried out as already described above.

A preferred embodiment of the method of the invention uses a purified alternansucrase. Purified alternansucrase is understood to mean an enzyme which is largely free from cell components of the cells in which the protein is synthesized and shows no contamination with proteins possessing polysaccharide-synthesizing activities (e.g. dextranases) or degrading activities, and/or no contamination with (polysaccharide) acceptors. The term "purified alternansucrase" preferably means an alternansucrase possessing a degree of purity of at least 70%, preferably at least 85%, and particularly preferably at least 95%.

The use of a purified protein for preparing alternan and/or fructose offers various advantages. Compared to methods working with partially purified protein extracts,

The reaction medium of the method of the invention does not contain any residues of the production strain (microorganism) which is used for the purification of the protein or for its preparation by genetic engineering.

Moreover, the use of the purified protein is advantageous for food and pharmaceutical industry applications. Thanks to the fact that the reaction medium is defined in its composition and freed from all unnecessary components, the product is likewise more precisely defined in respect of its components. In consequence of this, the procedure for obtaining food and pharmaceutical industry approval of these products produced by genetic engineering requires substantially less documentation, especially since these products should not show any traces of a transgenic microorganism.

Moreover, contrary to the so far described *in vitro* methods in cell free systems using partially purified alternansucrase preparations, the method of the invention using a purified alternansucrase has the advantage that it allows highly pure alternan to be prepared without the occurrence of dextransucrase and dextran contaminations, because of the high purity of the protein of the invention. Moreover, the method of the invention permits the production of alternan in high yields, without losses caused for instance by adverse side reactions of a dextransucrase, which would convert part of the substrate saccharose into undesired dextran, the separation of which from alternan would only be possible using time-consuming and expensive methods.

The method of the invention produces fructose in addition to alternan. The fructose can be used for the inexpensive recovery of so-called "high-fructose-containing syrups" (HFCS). The method of the invention yields products of high purity, because of the use of a purified alternansucrase. Hence, compared to conventional methods for preparing HFCS from maize starch, which comprise costly process steps for removing the buffer salts by ion exchange, (Crabb and Mitchinson, TIBTECH 15 (1997), 349-352) the method of the invention does not require an expensive purification of the fructose.

Another preferred embodiment of the method of the invention uses a recombinantly prepared alternansucrase.

According to another preferred embodiment, the enzyme possessing the enzymatic activity of an alternansucrase is immobilized on a carrier material.

Immobilization of the alternansucrase offers the advantage that the enzyme being the catalyst of the synthesis reaction can be easily recovered from the reaction mixture and reused several times. As the purification of enzymes is normally costly and time consuming, immobilization and reutilization of the enzymes allow for a substantial cost saving. Another advantage is the degree of purity of the reaction products not containing any residual protein.

There are many carrier materials available for the immobilization of proteins, and coupling to the carrier material can be made via covalent or non-covalent bonds (for an overview see: *Methods in Enzymology* 135, 136, 137). Widely used carrier materials include for instance agarose, alginate, cellulose, polyacrylamide, silica or nylon.

According to another embodiment of the invention, the alternansucrase (immobilized on a carrier material) is present between two membranes, one of which allows fructose, but not saccharose and alternan to penetrate, the other one of which allows saccharose, but not alternan to penetrate. The supply with substrate occurs through the membrane which allows saccharose to penetrate it. The synthesized alternan remains in the space between the two membranes and the released fructose can be continuously removed from the reaction equilibrium via the membrane which only allows fructose to penetrate it. Such an arrangement permits an efficient separation of the reaction products, and thus the production of pure fructose.

Moreover, the separation of fructose by ion exchange chromatography has been described ("*Starch Hydrolysis Products, Worldwide Technology, Production, and Application*", Edited by F. W. Schenck, R. E. Hebeda, (1992), VCH Publishers, Inc., New York).

Thus, the use of alternansucrases for preparing pure fructose on the one hand involves the advantage that the relatively inexpensive substrate saccharose can be used as the starting material, and on the other hand the fructose can be isolated easily from the reaction mixture without additional enzymatic conversions or chromatographic methods.

Moreover, the invention relates to methods for preparing alternan and/or fructose, wherein

- a) a saccharose-containing solution is contacted with a protein of the invention and acceptor molecules under conditions permitting the conversion of saccharose to alternan and/or fructose; and
- b) alternan and/or fructose is/are isolated from the solution.

Within the framework of the present invention an acceptor molecule is understood to mean a molecule at which an alternansucrase is able to catalyze a chain-extending reaction. The acceptor which can be added to the reaction mixture at the beginning of the reaction is preferably a carbohydrate or a carbohydrate derivative. The use of external acceptors leads to the production of low molecular products which are to be designated alternan in the context of the present invention. The carbohydrate acceptor is preferably an oligo or polysaccharide, in particular a branched polysaccharide, such as dextrin, glycogen or amylopectin, preferably a linear polysaccharide, and particularly preferably a saccharide selected from the group consisting of maltose, isomaltose, isomaltotriose and methyl- $\alpha$ -D-glucan. If an extension of the alternan chain at these acceptors occurs, then products are formed which have a higher molecular weight than the educt. Where maltose, isomaltose, isomaltotriose and methyl- $\alpha$ -D-glucan are used, one obtains products which have a lower molecular weight than the alternan that can be prepared in the absence of external carbohydrate acceptors.

The size of the molecular weight of the oligoalternans prepared depends on the saccharose/acceptor ratio used. For instance the degree of polymerization of the products increases as the saccharose/isomaltose ratio increases.

Moreover, the saccharose/acceptor ratio has an influence on the oligoalternan yield. For instance, the oligoalternan yield increases as the saccharose/isomaltose ratio decreases.

The hitherto described methods for producing oligoalternan with the use of alternansucrases which the authors claim have been purified (Pelenc et al., Sciences Des Aliments 11 (1991), 465-476) only yielded product mixtures of oligoalternan and oligodextran, in the presence of the carbohydrate acceptor maltose. In this case, the synthesis of oligodextran is presumably attributable to dextransucrase-contaminations of the alternansucrase preparation. Compared to

this method, the method of the invention offers the advantage that the use of recombinantly produced alternansucrase protein not containing any dextransucrase contaminants permits the preparation of oligoalternan without the simultaneous formation of oligodextran. Thus, the method of the invention makes it possible to provide oligoalternan, without requiring additional costly purification steps for separating oligodextran.

According to another preferred embodiment, the enzyme possessing the enzymatic activity of an alternansucrase is immobilized on a carrier material.

According to another preferred embodiment of the method of the invention, a recombinantly produced alternansucrase is used.

Moreover, the present invention relates to end products containing alternan. In this context, end products are understood to mean cosmetic products, preferably food products, fodder and particularly preferably pharmaceutical products.

Finally, the present invention relates to a method for preparing the afore-mentioned products comprising one of the above-described alternan manufacturing methods of the invention and the formulation of the thus obtained alternan in a form which is suitable for one of the afore-mentioned applications of the corresponding product.

These and other embodiments are disclosed and obvious to a skilled person and embraced by the description and the examples of the present invention. Additional literature regarding one of the above-mentioned methods, means and applications, which can be used within the meaning of the present invention, can be obtained from the state of the art, for instance from public libraries for instance by the use of electronic means. This purpose can be served inter alia by public databases, such as the "medline", which are accessible via internet, for instance under the address <http://www.ncbi.nlm.nih.gov/PubMed/medline.html>. Other databases and addresses are known to a skilled person and can be obtained from the internet, for instance under the address <http://www.lycos.com>. An overview of sources and information

regarding patents and patent applications in biotechnology is contained in Berks, TIBTECH 12 (1994), 352-364.

Description of the Figures:

Fig. 1:

Linear map of the entire sequence region which was cloned after the screening of a genomic library of *Leuconostoc mesenteroides* NRRL B 1355 by the corresponding overlapping fragments of the clones AS-19B1, AS-19B2, AS-28B and AS-29Ba.

Fig. 2:

Plasmid map pAlsu-pSK

Fig. 3:

HPLC chromatogram: Preparation of oligoalternan in the presence of maltose (Example 2).

Fig. 4:

Plasmid map pAlsu-pET24a

Fig. 5:

SDS PAGE with subsequent assay of sucrase activity (see Example 6)

The following protein extracts are used

1 + 2) *E. coli* BL21(DE3) containing pAlsu-pET24a-3

3 + 4) *E. coli* BL21(DE3) containing pAlsu-pET24a-7

5 + 6) *E. coli* BL21 (DE3) containing pAlsu-pET24a-21

7 + 8) *E. coli* BL21 (DE3) containing pET24a

1, 3, 5, 7) culture before induction with IPTG

2, 4, 6, 8) culture at the end of culturing

Fig. 6:

HPLC chromatogram of dextran T10

Fig. 7:

HPLC chromatogram of dextran T10 after dextranase digestion

Fig. 8:

HPLC chromatogram of oligoalternan

Fig. 9

HPLC chromatogram of oligoalternan after dextranase digestion.

Fig. 10

Map of the expression cassette including the polylinker of the plasmid pBinAR-N.

Fig. 11

Plasmid map pat-Alsu-Hyg.

Fig. 12

Plasmid map fnr-Alsu-Hyg.

### Examples

Vectors used in the examples:

#### 1. pBinAR-N

By the use of standard methods (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) we introduced a different polylinker (see figure 10) between the 35S Promoter and the OCS-Terminator into the plasmid pBinAR (Höfgen und Willmitzer, Plant Science 66 (1990), 221-230). The resulting plasmid was called pBinAR-N.

#### 2. pBinAR-Hyg-N

Via standard methods (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) we isolated an EcoRI/HinDIII-fragment from pBinAR-N containing the 35S Promoter, the polylinker and the OCS-Terminator. This fragment was then ligated into the same restriction

sites of plasmid pBIB-Hyg (Becker, Nucleic Acids Research 18 (1990), 203). The resulting plasmid was called pBinAR-Hyg-N.

### 3. pBinAR-pat-Hyg

By using the oligonucleotides Sp-pat-5' and Sp-pat-3' (s. SEQ ID Nos. 48 and SEQ ID No. 49) we amplified DNA molecules coding for the leader peptide of the patatin protein from potato (see SEQ ID No. 50, which differs from the sequence used by Sonnewald et al. Plant J. 1 (1991), 95-106 ) via a PCR approach using plasmid pgT5 (Rosahl et al., Mol. Gen. Genet. 203 (1986), 214-220; Sonnewald et al., Plant J. 1 (1991), 95-106) as a template. The resulting PCR products were cut by the restriction enzymes XbaI and Sall and then ligated into the plasmid pBinAR-Hyg-N which was linearized before by using the restriction enzymes SpeI and Sall. The resulting plasmid was called pBinAR-pat-Hyg.

PCR conditions:

Buffer and polymerase from Boehringer Mannheim (Pwo Polymerase No. 1644947)

DNA	0,2 ng
10x Buffer + MgSO <sub>4</sub>	5 µl
dNTPs (je 10 mM)	1 µl
Primer Sp-pat-5'	120 nM
Primer Sp-pat-3'	120 nM
Pwo Polymerase	1,0 units
distilled water	ad 50 µl

Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	64°C	0:30 min
Step 4	72°C	0:30 min
	(plus 1 sec per cycle)	
Step 5	72°C	5:00 min.

The steps 2 to 4 were repeated 35 times in a cyclical manner.

### 4. pBinAR-FNR-Hyg



By using the oligonucleotides Sp-fnr-5' and Sp-fnr-3 (see SEQ ID No. 51 and 52) we amplified DNA molecules coding for the transit peptide of the FNR protein from spinach via a PCR approach using plasmid p6SocFNR-15 (Jansen et al., Current Genetics 13, (1988), 517-522) as a template. The resulting PCR products were cut by XbaI and Sall and then cloned into the SpeI/Sall-opened pBinAR-Hyg-N. Resulting plasmid was called pBinAR-fnr-Hyg.

PCR conditions:

Buffer and polymerase from Gibco BRL (Platinum Taq DNA Polymerase High Fidelity No. 1304-011)

DNA	0,2 ng
10x Buffer	5 µl
MgSO <sub>4</sub>	2,0 µl
dNTPs (per 10mM)	1 µl
Primer Sp-fnr-5'	150 nM
Primer Sp-fnr-3'	150 nM
Taq Platinum Hifi Polymerase	1,5 units
distilled water	ad 50 µl

Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	58°C	0:30 min
Step 4	68°C	0:20 min
	(plus 1 sec per cycle)	
Step 5	68°C	3:00 min

The steps 2 to 4 were repeated 35 times in a cyclical manner.

**Example 1: Cloning of alternansucrase from *Leuconostoc mesenteroides* NRRL-B1355**Isolation and sequencing of alternansucrase

The strain *Leuconostoc mesenteroides* NRRL-B1355 was cultured in 1 l of Lactobacilli MRS Broth (Difco) complemented with 5% saccharose at 28°C for two days. After the culture was subjected to centrifugation at 20,000 x g for 30 minutes, the supernatant was admixed with the same volume of 10% trichloro acetic acid and stirred at 4°C for 16 hours. This solution was then subjected to centrifugation at 10,000 x g for 30 minutes. The thus obtained precipitate was dissolved in 4.5 ml of 40 mM Tris-HCl, pH 8.8, and subsequently neutralized with (about 0.5 ml) 2 M Tris-base. This protein solution was given to the company Toplab Gesellschaft für angewandte Biotechnologie mbH, Martinsried, Germany, for protein sequencing. At this company, the protein solution was electrophoretically separated in SDS polyacrylamide gel, the gel was stained with Coomassie Blue and the staining was subsequently removed by 10% acetic acid. For the enzymatic digestion of the protein, the protein bands were cut from the gel, pressed through a sieve and fragmented (pores 30 µm x 100 µm). The crushed gel was then washed with half concentrated incubation buffer (12.5 mM Tris, 0.5 mM EDTA pH 8.5) for 2 minutes. Subsequently, it was subjected to centrifugation, the buffer was removed and the gel was dried in the "Speedvac" for one hour (about 5% residual water, rubber-like). Subsequently, a solution of endoproteinase LysC in 400 µl 12.5 mM Tris/HCl, pH 8.5 (enzyme: protein = 1 : 10) and 0.1% of laurylmaltoside was prepared. 200 µl of this solution were added to the sample and incubated in the heat block shaker at 37°C overnight. In order to elute the peptide fragments, a one hour incubation with 1% TFA was carried out, twice, followed by centrifugation, and subsequently by elution with 10% formic acid, 20% isopropanol, 60% acetonitrile for 3 hours. The peptide fragments obtained were then separated from each other by HPLC (column Superspher 60 RP select B (Merck, Darmstadt) 2 mm x 125 mm; buffer A 0.1% trifluoro acetic acid, buffer B: 0.085% TFA in acetonitrile; flow rate: 0.2 ml/min; gradient: 5-60% in 60 min; detection at 206 nm. The peptide fragments obtained were then sequenced in an automatic sequencer Procise 492 (Applied Biosystems,

PE); the procedure being the stepwise Edman degradation in a modification according to Hunkapiller (Hunkapiller et al., Meth. Enzymol. 91 (1983), 399-413).

Six different peptide sequences (see Seq. ID Nos. 5 to 9, Seq. ID No. 21) were identified which were designated lysC-66, lysC-67, lysC-82, lysC-83, lysC-88 and "N-terminus".

#### Preparation of a genomic DNA library from *Leuconostoc mesenteroides* NRRL B1355

*Leuconostoc mesenteroides* NRRL-B1355 (purchased from ATCC) was cultured in 100 ml YT medium (Sambrook et al, loc. cit.) additionally containing 2% (w/v) of glucose and 50 mM sodium phosphate buffer pH 7.0, at 28°C for 36 hours. After harvesting the cells by centrifugation, genomic DNA was isolated according to Ausubel et al. (Current Protocols in Molecular Biology, Volume 1, Greene and John Wiley & Sons (1994), USA).

100 µg of genomic DNA from *Leuconostoc mesenteroides* NRRL-B1355 were partially digested with 0.001 units of the restriction enzyme Sau3A for 30 minutes, subsequently extracted with phenol:chloroform:isoamyl alcohol (25:24:1) and precipitated with ethanol. 2.5 µg of the partially digested DNA obtained from *Leuconostoc mesenteroides* NRRL-B1355 were ligated with T4 DNA ligase in 1 µg of the BamHI-cut and dephosphorylated vector pBKCMVBamHI (Stratagene) under the conditions indicated by the manufacturer (Stratagene, pBK phagemid vectors instruction manual & T4 DNA ligase ligation kit). 2 µl of the ligation mixture were packaged with Gigapack III Gold (Stratagene) according to the instructions of the manufacturer and stored after the amount of phage content had been determined.

#### Preparation of the probe for isolating the alternansucrase gene

From the peptide sequences lysC-66 (Seq. ID No. 5), lysC-67 (Seq. ID No. 6), lysC-82 (Seq. ID No. 7), lysC-83 (Seq. ID No. 8) and lysC-88 (Seq. ID No. 9) obtained after tryptic digestion of the purified alternansucrase protein (see above) the peptides lysC-82 and lysC-83, after having undergone reverse translation, were selected for the synthesis of degenerated oligonucleotides (Seq. ID No. 10, Seq. ID No. 11). Said oligonucleotides served as primers in a PCR reaction on genomic

DNA of NRRL-B1355. All positions within oligonucleotides depicted as N were replaced by inosin in the primer synthesis.

#### PCR Reaction conditions

The reaction mixture was prepared with the buffers supplied for Taq polymerase (Company GibcoBRL).

Reaction mixture:

#### Taq Polymerase (Gibco)

DNA	100	ng (genomic NRRL-B1355)
DNTPs	2.5	mM for each nucleotide
primer	10	μl of a solution containing 0.2 μMol
10 fold buffer	5	μl
magnesium chloride	2	mM
polymerase	1	unit
water	ad	50 μl

Step 1	95°C	3'
Step 2	95°C	1'
Step 3	58°C	2'
Step 4	72°C	2'
Step 5	72°C	10'

40 repetitions of steps 2 to 4

An 837 bp fragment (Seq. ID No. 12) resulting from this PCR reaction, the ends of which were blunted with T4 DNA polymerase, was cloned into the SmaI-cut pBlueSkript vector (Stratagene). The resulting plasmid was designated pAlsu-PCR-lysc82/83. After sequencing of the insert and computer-aided translation into the corresponding protein sequences, a data base comparison was carried out in the Swiss Prot data base. This comparison showed homologies to known glycosyl transferases (P49331, P11001, P68987, P13470, P27470, P29336).

About 5,000 phages of the genomic DNA library of *Leuconostoc mesenteroides* NRRL-B1355 were plated out using the bacterial strains and nutrient solutions indicated by the manufacturer (Stratagene), and after incubation at 37°C for 12 hours were transferred to nitrocellulose filters. This was followed by denaturation of the phages by immersion of the nitrocellulose filters in 1.5 M sodium chloride, 0.5 M caustic soda solution for 2 minutes and neutralization of the filters by immersion in 1.5 M sodium chloride, 0.5 M Tris-HCl, pH 8.0 for 5 minutes. After rinsing the filters in 0.2 M Tris-HCl, 2 x SSC, the phage DNA was bound to the membranes by UV cross link (Stratalinker of the company Stratagene, 120,000  $\mu$ J for 30 seconds). The filters were incubated in a prehybridization solution (5 x SSC, 0.5% BSA, 5 x Denhardt, 1% SDS, 40 mM sodium phosphate buffer, pH 7.2, 100 mg/l herring sperm-DNA, 25% formamide) at 42°C for 6 hours. 30 ng of the isolated insert from the plasmid pAlsu-PCR-lysc82/83 were radioactively labeled by means of a multiprime kit (Boehringer Mannheim) using  $\alpha$ -  $^{32}$ P dCTP (ICN Biomedicals). This radioactive probe was added to the prehybridization mixture and the filters were incubated in this hybridization mixture at 42°C overnight. After removal of the hybridization mixture the filters were washed three times in a washing solution (0.1 x SSC, 0.5% SDS) at 55°C for 15 minutes. An X-ray film (Kodak) was then placed on the filter for 18 hours. Phage colonies, producing hybridization signals, were identified, isolated, resuspended in SM medium and then again plated out in a dissolution such that they could be recognized as single plaques. After these phages were transferred to nitrocellulose filters and subjected to further treatment and hybridization under conditions as described above, hybridizing phages were obtained as individual isolates by means of the radioactive gene probe used. After *in vivo* excision of the isolated phages in accordance with the manufacturer's instructions (Stratagene) the clones AS-19B1 and AS-19B2 could be isolated as plasmids. After complete sequencing of both clones (Agowa) (Seq. ID No. 13, Seq. ID No. 14) both sequences showed an 1008 bp overlap. The joining of Seq. ID No. 13 with Seq. No. 14 followed by computer aided translation of all possible reading frames allowed a continuous reading frame, starting with the codon ATG (corresponding to the bases 678 to 680 in Seq. ID No. 1), to be identified. As no stop codon could be found in this composed reading frame, additional clones were isolated in order to obtain the complete coding sequence of alternansucrase.

Therefore, about 5,000 phages of the genomic DNA library of *L. mesenteroides* NRRL-B1355 were again examined for hybridization by means of a clone AS-19B2 subfragment radioactively labeled using the multiprime kit (Boehringer Mannheim), as described above. The hybridization probe was prepared with the use of the HindIII (restriction site in the insert of AS-19B2) / Sall (cuts the pBKCMV phagemid vector in the polylinker)-fragment from AS-19B2. Said fragment contains 372 bases of the 3' end of the sequences encoding the above-described reading frame. The screening of the phage library, singling out, and transformation of the phages into plasmids was carried out under the above-described conditions. After complete sequence analysis of the thus isolated clones AS-28B (see Seq. ID No. 15) and AS-29Ba (Seq. ID No. 16) it was possible to identify an overlap of 960 identical bases (corresponding to bases 4863 to 5823 in Seq. ID No. 1) between clones AS-19B2 (Seq. ID No. 14) and AS-28B and an overlap of 567 identical bases (corresponding to bases 5256 to 5823 in Seq. ID No. 1) between clones AS-19B2 and AS-29Ba (Seq. ID No. 16). Clones AS-28B and AS-29Ba have 1523 identical bases (corresponding to bases 5256 to 6779 in Seq. ID No. 1). After computer-aided joining of clones AS-19B1, AS-19B2 and AS-28B a continuous reading frame starting with codon ATG (bases 678 to 680 on the complete sequence) appeared. This reading frame also does not contain a stop codon. After the joining of clones AS-19B1, AS-19B2, AS-28B and AS-29Ba it was possible to identify a reading frame starting with the codon "ATG" (corresponding to bases 678 to 680 in Seq. ID No. 1) and ending with "TAA" (corresponding to bases 6849 to 6851 in Seq. ID No. 1) encoding 2057 amino acids. In addition to the coding region, the entire isolated and identified DNA sequence of the composed clones (Seq. ID Nos. 13 - 16) contains 677 bases in the 5' region and 2469 bases in the 3' region which represent sequences not encoding alternansucrase (see Fig. 1).

**Example 2: Construction of plasmid pAls-pSK for the transformation of *E. coli* and test of the protein extracts for enzymatic activity**

Plasmids AS-19B1, AS-19B2, AS-28B and AS-29Ba (see Example 1) were joined in the following manner: A NotI-(restriction site in the polylinker of vector pBK CMV, company Novagen)/Clal-fragment of clone AS-19B1 was inserted into the vector

pBluescript SK (company Stratagene) at the same restriction sites (= first cloning step). Consecutive insertion of the ClaI/XhoI fragment from AS-19B2, XhoI/MluI fragment from AS-28B and MluI/BsaBI (BsaBI-cut fragment cloned into the blunted Apal restriction site of the vector) fragment of AS-28B into the clone obtained from the first cloning step produced plasmid pAlsu-pSK (see Fig. 2). This plasmid contains the complete coding sequence of the alternansucrase from *Leuconostoc mesenteroides* NRRL-B1355 as well as non-coding sequences of 677 bp (promoter region) in the 5' region and 539 bp in the 3' region (Seq. ID No. 17).

Plasmid pAlsu-pSK was then transformed in *E. coli* (DH5 $\alpha$  company Fermentas). The bacteria were then cultured at 27°C for two days in 50 ml "Terrific broth" (the composition of which is described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2<sup>nd</sup> edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (supplemented with 0.5% glucose) or in a fermentation medium having the following composition: KH<sub>2</sub>PO<sub>4</sub> 1.5 g/l, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 5.0 g/l, NaCl 0.5 g/l, Na-citrate 1.0 g/l, Fe<sup>2+</sup>SO<sub>4</sub> x 7 H<sub>2</sub>O, 0.075 g/l, yeast extract 0.5 g/l, tryptone 1.0 g/l, glucose 15.0 g/l, MgSO<sub>4</sub> x 7, H<sub>2</sub>O 0.3 g/l, CaCl<sub>2</sub> x 2 H<sub>2</sub>O 0.014 g/l, mineral salts 10 ml/l, H<sub>3</sub>BO<sub>3</sub> 2.5 g/l, CoCl<sub>2</sub> x 6 H<sub>2</sub>O 0.7 g/l, CuSO<sub>4</sub> x 5 H<sub>2</sub>O 0.25 g/l, MnCl<sub>2</sub> x 4 H<sub>2</sub>O, 1.6 g/l, ZnSO<sub>4</sub> x 7 H<sub>2</sub>O 0.3 g/l, Na<sub>2</sub>MoO<sub>4</sub> x 2 H<sub>2</sub>O 0.15 g/l, vitamin B1 (thiamine) 0.005 g/l.

All cultures contained 100 mg/l ampicillin. The cells were then harvested by centrifugation, resuspended in 2 ml 50 mM Na-phosphate buffer pH 7.2 and crushed by a French Press. Subsequently, they were again subjected to centrifugation to remove solid particles of the crushed cells, and the supernatant (hereinafter referred to as (protein) extract) was used after sterilfiltration (Sterivex GV 0.2  $\mu$ m, millipore) for further analyses.

#### In vitro preparation of alternan by means of protein extracts

For the in vitro preparation of alternan, 200  $\mu$ l each of the extracts obtained were examined in 2 ml each of 100 mM Na-citrate buffer pH 6.5 and 20% (w/v) saccharose for activity in the presence and absence of 100  $\mu$ l of 10 mM maltose. The reaction mixture was incubated at 37°C for 24 hours. In the subsequent

precipitation with the same volume of ethanol in the absence of maltose no precipitable polymer was found. In the batch containing maltose, HPLC chromatography (Dionex PA-100 column, running buffer 150 mM NaOH, elution buffer 150 mM NaOH + 3 M sodium acetate buffer gradient) showed the formation of oligomers (see Fig. 3).

#### Activity gel

20 ml each of the individual protein extracts were applied to a 6% SDS-PAA gel and separated at a current strength of 20 mA per gel. (Before application to the gels, the extracts were not incubated at 95°C). Subsequently, the extracts were examined for sucrase activity according to the method of Miller and Robyt (Analytical Biochemistry 156 (1986), 357-363).

The control (dextransucrase NRRL-B-512F, see Example 3 for its preparation) showed polymerizing activity. The protein extracts of the above-described *E. coli* cells containing the plasmid pAIsu-pSK, did not show any polymer-forming activity.

### **Example 3: Cloning and expression of dextransucrase from *Leuconostoc mesenteroides* NRRL-B512F**

#### Isolation of genomic DNA

*Leuconostoc mesenteroides* NRRL-B512F (obtained from ATCC) was cultured at 28°C for 48 hours in YT-medium (Sambrook et al., Molecular Cloning: A Laboratory Course Manual, 2<sup>nd</sup> edition (1989), Cold Spring Harbor Press, New York) additionally containing 1% of saccharose and 50 mM sodium phosphate buffer pH 7.0. After harvesting the cells by centrifugation, genomic DNA was isolated according to Ausubel et al. (Current Protocols in Molecular Biology, Volume 1, Greene and John Wiley & Sons (1994), USA).

#### PCR Amplification of the Dextransucrase Gene and Cloning in pET24a

For the recombinant expression of dextransucrase in *E. coli*, the gene encoding dextransucrase was cloned in the expression vector pET24a (Novagen) after PCR amplification. For this purpose, an EagI restriction site was introduced at the 5' end of the sequences encoding the dextransucrase and an XhoI restriction site at the 3'



end, together with the PCR primers used (5'b512-1: 5'-ACTgCggCCgCATgCCATTTACAgAAAAg-3'; Seq. ID No. 3 and 3'b512: 5'-ACTgCTCgAgTTATgCTgACACAgCATTTTC-3'; Seq. ID No. 4) derived from the sequence of WO 89/12386. Subsequent cloning into the corresponding restriction sites of the polylinker of pET24a was carried out. The resulting plasmid was designated UL5-20.

#### PCR Reaction conditions

Buffer and polymerase of the company Gibco BRL were used.

DNA:	100	ng (genomic NRRL-B512F)
10 fold buffer	5	μl
MgCl <sub>2</sub>	4	mM
5' primer	50	ng
3' primer	50	ng
dNTP	1	mM of each nucleotide
Pfu polymerase	0.5	units
water	ad 50	μl

step 1	95°C	4 minutes
step 2	95°C	1 minute
step 3	55°C	1 minute
step 4	72°C	5 minutes
step 5	72°C	10 minutes

40 repetitions were made between steps 2 and 4.

#### Preparation of recombinant dextransucrase

BL21(DE3) *E. coli* cells containing the plasmid UL5-20 were cultured in YT medium (see above) at 37°C up to an OD<sub>600</sub> = 0.8. Subsequently, the cells were subjected to induction with 0.2 mM IPTG and cultured anew at 18°C for 24 hours. After harvesting the cells by centrifugation and resuspending them in sodium phosphate buffer, pH 5.2, the cells were crushed in a French Press. The solution obtained

was freed from insoluble components by centrifugation and the supernatant containing dextranucrase and referred to hereinafter as the extract was obtained.

#### **Example 4: PCR Amplification of the Coding Region of Alternansucrase and Cloning in pET24a**

The coding region of alternansucrase was amplified in a PCR reaction (see the reaction conditions below) with genomic DNA from the *Leuconostoc mesenteroides* strain NRRL-B1355 as a template. An NheI restriction site was introduced at the 5' end by means of primers A1-4 (Seq. ID No. 18), and a Sall-restriction site at the 3' end by means of primer A1-5 (Seq. ID No. 19). A fragment of about 6200 bp was isolated.

A1-4: 5'-GGG CCC GCT AGC ATG AAA CAA CAA GAA ACA GT

A1-5: 5'-CCC GGG GTC GAC CTT TGT CGA ATC CTT CCC

Reaction conditions of the PCR (kit of the company Gibco BRL):

DNA	1 µl
10 x buffer	5 µl
10 mM per dNTP	2 µl
50 mM MgSO <sub>4</sub>	2 µl
primer	per 1 µl
Platinum DNA polymerase	0.2 µl
distilled water	37.8 µl
step 1	95°C, 2 minutes
step 2	95°C, 20 seconds
step 3	47°C, 20 seconds
step 4	68°C, 7 minutes (prolonged by 3 seconds per cycle)
step 5	68°C, 15 minutes

Steps 2 to 4 were repeated 35 times altogether before step 5 was carried out.

The PCR fragment obtained was purified according to standard methods, treated with the restriction endonucleases NheI and SalI, ligated into vector pET24a (of the company Novagen) which had likewise been cut with these enzymes, and the ligation product was transformed into *E. coli*. After preparation of the plasmid and restriction digestion, three positive clones were selected. They were designated pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21 (see Fig. 4), respectively. All contained the sequence indicated in Seq. ID No. 20 as an insertion.

### **Example 5: Expression of the Recombinant Alternansucrase in *E. coli* in Shake Flask Cultures and in the Fermenter**

#### Shake Flask Culture

Plasmids pAlsu-pET24a-3, pAlsu-pET24a-7, pAlsu-pET24a-21 and pET24a were transformed into *E. coli* BL21 (DE3), of the company Novagen, and after initial culturing at 37°C for 3 hours in 3 ml YT medium (Sambrook et al., Molecular Cloning, A Laboratory Manual, 2<sup>nd</sup> edition (1989), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) they were each cultured in shake flasks in 2 replicas in 50 ml Davis minimal medium (DIFCO Manual, Dehydrated Culture Media and Reagents for Microbiology, 10<sup>th</sup> edition, Detroit Michigan, USA (1984)) containing 0.2% glucose instead of dextrose as a carbon source at 37°C until an OD<sub>600</sub> of about 0.8 was reached. After centrifugation and resuspension, one of the two replica cultures was cultured in Davis Minimal Medium (DMA) containing 1% lactose as the carbon source and inductor at 27°C for another 16 hours. The cells of the individual cultures were harvested after centrifugation, resuspended in 50 mM sodium acetate buffer pH 5.3, and a protein extract was prepared as described in Example 2.

#### Fermenter

Clone pAlsu-pET24a-21 transformed in *E. coli* BL21(DE3) was cultured in a 2 l fermenter (Biostad B; B.Braun, Melsungen) under the following conditions:

Medium:

Fermentation medium:  $\text{KH}_2\text{PO}_4$  1.5 g/l,  $(\text{NH}_4)_2\text{SO}_4$  5.0 g/l, NaCl 0.5 g/l, Na-citrate 1.0 g/l,  $\text{Fe}^{2+}\text{SO}_4 \times 7 \text{ H}_2\text{O}$  0.075 g/l, yeast extract 0.5 g/l, tryptone 1.0 g/l, glucose 15.0 g/l,  $\text{MgSO}_4 \times 7 \text{ H}_2\text{O}$  0.3 g/l,  $\text{CaCl}_2 \times 2 \text{ H}_2\text{O}$  0.014 g/l, mineral salts 10 ml/l,  $\text{H}_3\text{BO}_3$  2.5 g/l,  $\text{CoCl}_2 \times 6 \text{ H}_2\text{O}$  0.7 g/l,  $\text{CuSO}_4 \times 5 \text{ H}_2\text{O}$  0.25 g/l,  $\text{MnCl}_2 \times 4 \text{ H}_2\text{O}$  1.6 g/l,  $\text{ZnSO}_4 \times 7 \text{ H}_2\text{O}$  0.3 g/l,  $\text{Na}_2\text{MoO}_4 \times 2 \text{ H}_2\text{O}$  0.15 g/l, vitamin B1 (thiamine) 0.005 g/l.

Carbon source: Glucose (1.5% (w/v)) is present in the medium, 70% (w/v) glucose solution is added.

Automatic pH control by ammonia and phosphoric acid at pH 7.0  $\pm$  0.1. A 20% concentration of  $\text{pO}_2$  is adjusted in the medium via control by the stirrer.

Conditions:

1.5 l of fermentation medium were inoculated with 50 ml of the preculture.

The cells were first cultured at 37°C until the glucose present was consumed. They were then cultured at the same temperature at a feeding rate of 9 g of glucose  $\times \text{l}^{-1} \times \text{h}^{-1}$  until an  $\text{OD}_{600} = 40$  was reached. At this time, the temperature of the culture broth was lowered to 20°C and the amount of glucose addition was lowered to 2 g  $\times \text{l}^{-2} \times \text{h}^{-1}$ . At a culture temperature of 20°C, the culture was subjected to induction with 0.2 mM IPTG (isopropyl- $\beta$ -D-thiogalactopyranoside (Sigma)). After culturing at 20°C for another 18 hours, the cells were harvested by centrifugation, resuspended in 50 mM sodium phosphate buffer pH 5.3 and an extract was prepared as described in Example 2.

#### **Example 6: SDS PAGE Assay of the Activity of the Recombinant Alternansucrase, Periodic Acid Oxidation and Staining according to Schiff**

Protein extracts were prepared from *E. coli* shake flask cultures (strain BL21 (DE3)), containing the plasmids pAlsu-pET24a-3, pAlsu-pET24a-7, pAlsu-pET24a-21 and pET24a (control), respectively. Two different extracts were each prepared from the cells transformed with the different extracts, one of said extracts being prepared before induction with IPTG and the other one being prepared after induction with IPTG at the end of culturing. The activity of these extracts of shake

flask cultures (see Example 5) was detected by SDS PAGE separation of the proteins, followed by SDS removal by washing with 50 mM sodium acetate buffer pH 5.3 and incubation of the gels in 50 mM sodium acetate pH 5.3, 5% (w/v) saccharose at 37°C for 16 hours, followed by periodic acid oxidation of the polymer formed and staining by means of acidic Schiff reagent (Miller and Robyt, Analytical Biochemistry 156, (1986), 357-363).

Fig. 5 shows that sucrase activity has not been found for either one of the extracts (preparation of the extract before and after IPTG-induction) containing the cloning vector pET24a. In the case of strains which had been transformed with the plasmids pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21, respectively, all protein extracts showed sucrase activity at the end of the induction phase (concentrated in one band).

Before induction with IPTG such activity bands were not found.

As the polymer formed in the gel can be stained according to the above-described methods by acidic Schiff reagent, it can be assumed not to be composed of pure  $\alpha$ -1,3-linked units which would not lead to any staining.

As the gene contained in vectors pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21, respectively, was isolated from the *Leuconostoc mesenteroides* strain NRRL-B1355 which expresses at least one dextran sucrase apart from alternansucrase, it was not possible to determine unambiguously with this staining method whether the nucleic acid sequence contained in the plasmid actually encodes an alternansucrase. Dextrans and alternans can both be detected by this method because both polymers contain  $\alpha$ -1,6 linkages.

#### **Example 7: Tests for the Enzymatic Activity of Recombinantly Prepared Alternansucrases after Heat Treatment and for the Specificity of Alternansucrase**

In order to prove polymerization activities, extracts from shake flask cultures were used (see Example 5). 100  $\mu$ l of extract were each added to 2 ml reaction buffer (50 mM sodium acetate pH 5.3, 20% saccharose) and incubated at 37°C for 24 hours. For comparison, an extract inactivated by a 10 minute treatment at 95°C, and an

extract from *E. coli* BL21(DE3) containing vector pET24a were used. Polymer formation was only found in the batch that had not been inactivated, while the batch treated at 95°C for 10 minutes and the batch with the extract from BL21(DE3) containing pET24a did not show any polymer formation. After addition of the same volume of absolute ethanol to all batches, polymers could only be precipitated from the batch which had not been inactivated. This finding is a clear indication of the activity of alternansucrase, because the dextranucrase present in NRRL B-1355 is inactivated by a treatment at 45°C for 30 minutes, while alternansucrase remains active under these conditions (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85). The enzymatic assay by a coupled enzymatic test of the glucose and fructose released and of the saccharose still contained in the reaction mixture after 24 hours, respectively, revealed that fructose was only present in the extract that was not inactivated.

For carrying out the enzymatic test either purified protein or crude protein extract is added in different dilutions to 1 ml batches containing 5% saccharose and 50 mM acetate, pH 5.5 and subjected to incubation at 37°C. After 5 minutes, 10 minutes, 15 minutes, 20 minutes, 25 minutes and 30 minutes, 10 µl each are removed from these batches and the enzymatic activity of alternansucrase is terminated by immediate heating to 95°C. Subsequently, in the coupled photometric test, the portions of fructose and glucose released by alternansucrase and the portion of used-up saccharose, respectively, are determined. For this purpose, 1 µl to 10 µl of the inactivated sample are placed into 1 ml of 50 mM imidazole buffer, pH 6.9, 2 mM MgCl<sub>2</sub>, 1 mM ATP, 0.4 mM NAD and 0.5 U/ml hexokinase. After sequential addition of about 1 u of glucose-6-phosphate dehydrogenase (from *Leuconostoc mesenteroides*), about 1 u of phosphoglucose isomerase and about 5 u of invertase, the alteration of adsorption at 340 nm is measured. Subsequently, the amount of fructose and glucose released and used-up saccharose, respectively, is calculated according to the Lambert-Beer law.

In control batches (inactivation of the extract by treatment with 95°C and extract from *E. coli* containing pET24a) no significant release of fructose and no decrease of saccharose, respectively, was found in the reaction batch after 24 hours.

These results confirm that the specificity of the sucrase encoded by plasmids pAIsu-pET24a-3, pAIsu-pET24a-7 and pAIsu-pET24a-21, respectively, is that of a

glucosyltransferase. The specificity of a fructosyl transferase, the presence of which has been described for some strains of the genus *Leuconostoc* is to be excluded, because otherwise glucose should have been found.

**Example 8: Production of Alternan by Means of Alternansucrase Prepared in *E. coli***

100 ml of extract obtained by fermentation of *E. coli* BL21(DE3) containing plasmid pAlsu-pET24a-3 (see Example 4) were added to 900 ml of reaction buffer (50 mM sodium acetate pH 5.3, 20% saccharose) and incubated at 37°C for 24 hours. The addition of the same amount of absolute ethanol to the reaction mixture caused the alternan formed to precipitate. After the precipitate was washed twice with 50% ethanol, it was dried by lyophilization. The yield of dried polymer based on the amount of saccharose used in the reaction was 60%.

**Example 9: HPLC Analysis of Alternan and Dextran After Dextranase Digestion**

100 mg of the polymer prepared in Example 7 and 100 mg of dextran T10 (Pharmacia) were each dissolved in 1 ml of water. 40 µl each of these solutions were added to 700 µl reaction buffer (50 mM potassium phosphate pH 5.7, 8 units of dextranase, ICN Biomedicals Inc. No. 190097), and incubated at 37°C for 16 hours. 50 µl of the polymer solutions not treated with dextranase (see Fig. 6) and 50 µl of the polymer solutions treated with dextranase (Fig. 7) were analyzed by HPLC (Dionex, column PA-100, NaOH / NaOH-NaAc gradient).

In the case of dextran T10 the cleavage of the polymer into different molecules of lower molecular weights can be clearly seen. The entire high molecular weight dextran is converted by dextranase into smaller units (mostly isomaltose). By contrast, in the case of alternan, short chained oligosaccharides only appear in small amounts after dextranase incubation. Most of the alternan is not digestible by dextranase. This finding suggests that the product prepared by recombinant alternansucrase is not dextran, but alternan which is known to be hardly accessible

to enzymatic digestion by dextranase (Lopez-Mungia et al., Enzyme Microb. Technol. 15, (1993), 77-85).

#### **Example 10: *In vitro* Preparation of Alternan in the Absence of Dextranase**

100  $\mu$ l extract from shake flask cultures (see Example 5) were added to 2 ml of reaction buffer (50 mM sodium acetate, pH 5.3, 20% saccharose). 50 units of dextranase (Biomedicals Inc. No. 190097) were additionally added to another batch. Two corresponding batches which contained dextransucrase from *Leuconostoc mesenteroides* NRRL-B512F instead of the enzyme extract served as controls; one of these two batches had dextranase additionally admixed to it.

After precipitation with ethanol, the reaction batch with dextransucrase and dextranase did not show any polymer formation. All other batches were found to show polymer formation.

#### **Example 11: *In vitro* preparation of Oligoalternan and HPLC Analysis**

Oligoalternan was prepared as described in Example 2, with a protein extract in the presence of maltose and was subsequently detected (See Fig. 8) by HPLC-chromatography (see Example 2). For comparison, a portion of this batch was admixed with 50 units of dextranase (Biomedicals Inc. 190097) after preparation of oligoalternan and subsequently separation by HPLC chromatography was carried out as well (see Fig. 9). A comparison of the two chromatograms shows that not only the height of the two peaks which can be allocated to the oligoalternan ( $\alpha$  and  $\beta$ -anomer) (retention time between 15.87 and 16.61 minutes) but also the height of all the other peaks, the first signs of which are already visible without dextranase, remain unchanged. This finding suggests that recombinantly prepared alternansucrase allows oligoalternan to be prepared without the simultaneous production of oligodextran. Oligodextran would be liable to digestion by dextranase, which would have to show up in a decrease of the height of the peaks in the HPLC chromatogram, if oligodextran were present.



## Example 12: Methylation analysis of Alternan

In order to further analyze the alternan produced *in vitro* a methylation analysis was carried out:

### Permethylation

The permethylation was performed as described by Ciucanu and Kerek (Carbohydr. Res. 131 (1984), 209-218) by using NaOH/Mel in DMSO or by using a modified method according to Hakomori (Journal of Biochemistry 55 (1964 FEB), 205-208) which relies on the use of freshly prepared Li-Dimsyl/Mel (Dimsyl= methylsulfinyl carbanion) in DMSO at room temperature.

All reactions are performed under a nitrogen atmosphere. The permethylation products are isolated by extracting the excess of methyl iodide by the use of dichloromethane. DMSO and salts were washed out at the end.

### Degradation into partially methylated sorbitacetates (methylation analysis)

The permethylated glucans were hydrolyzed with 2N trifluoroacetic acid at 120°C for 1-3 hours. After cooling the acid was removed by nitrogen. Then the resulting glucans were co-distilled with a small amount of toluene, afterwards reduced by NaBD<sub>4</sub> in 1N ammonia and finally, acetylated by pyridine/acetic anhydride (3h, 90°C). The products were extracted by dichloromethane and washed with NaHCO<sub>3</sub>. The products in the organic phase were analyzed by gas chromatography.

### Analysis of the acetylated products

The acetylated products were analyzed by gas chromatography which was performed with a chromatograph manufactured by the Carlo-Erba company model GC 6000 Vega equipped with an on-column injector, a 25m CPSol8CB and a FID-detector. As a carrier gas hydrogen (80 kPa) was used.

The identification and integration of the peaks was performed as described by Sweet et al. (Carbohydr. Res. 40 (1975), 217).

### Results

The following main components were identified by gas chromatography:

Sorbit acetylated in position	Interpretation
1,5	Terminal Glucopyranose
1, 3, 5	3-linked Glucopyranose
1, 5, 6	6-linked Glucopyranose
1, 3, 5, 6	3,6-linked Glucopyranose

Furthermore, small amounts (rel. amount 0.2-0.4 mol%) of the following components were also found: 1, 4, 5- and 1, 3, 4, 5-sorbit and another tetraacetyl component (1,5,x,y). It is supposed that these components are due to incomplete methylation.

The following amounts were found for the above mentioned components in different experiments which were performed by changing the length of hydrolysis (indicated in bold by the number of hours) (MA=methylation analysis1; MA-b= methylation analysis 2):

Values in mol%

Ac in Pos	MA (1h)	MA (2h)	MA (3h)	MA-b(2h)
1,5	10,49	10,56	9,17	12,71
1,3,5	31,69	34,70	32,95	23,12
1,4,5	0,70	0,30	0,36	0,33
1,5,6	47,02	44,17	47,23	54,62
1,3,4,5	0,27	0,22	0,25	0,31
1,5,x,y	0,19	0,32	0,36	0,24
1,3,5,6	9,64	9,73	9,68	8,67

**Example 13: Construction of an expression cassette for plants: vacuolar and plastidic expression of an alternansucrase.**

By using plasmid Als-u-pET24a as a template and the PCR primers Al-5'-1.2 and Al-3'-2.2 (see SEQ ID NO 53 and 54) we amplified the coding region of alternansucrase from *Leuconostoc mesenteroides* which was then cut by the

restriction enzymes Sall and PstI. Afterwards the resulting fragments were cloned into Sall and SdaI digested plasmids a) pBinAR-pat-Hyg and b) pBinAR-fnr-Hyg.

The resulting plasmids were called a) pat-Alsu-Hyg (see figure 11) and b) fnr-Alsu-Hyg (see figure 12).

Note: The bacterial secretion signal peptide was removed from the cds by choice of the PCR primers.

#### PCR conditions:

Buffer and polymerase from Boehringer Mannheim (Pwo Polymerase No. 1644947)

DNA	0,5 ng
10x Buffer + MgSO <sub>4</sub>	5 µl
dNTPs (je 10 mM)	2 µl
Primer Sp-AS-5'	100 nM
Primer Sp-AS-3'	100 nM
Pwo Polymerase	1,0 unit
distilled water	ad 50 µl

#### Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	47°C	0:30 min
Step 4	68°C	7:00 min
	(plus 3 sec per cycle)	
Step 5	68°C	15:00 min

The steps 2 to 4 were repeated 35 times in a cyclical manner.

#### **Example 14: Northern blot analysis for expression of alternansucrase in transgenic plants**

Leaves or tubers from potato plants transformed via agrobacteria with plasmids pat-Alsu-Hyg and fnr-Alsu-Hyg, respectively, were pulverized in a mill, type MM 200, (Retsch GmbH & Co. KG, 42781 Haan, Germany) at 30 Hz for 50 sec. RNA was extracted according to Logemann et al. (Anal. Biochem. 163 (1987), 16-20). 50 µg

RNA per sample were loaded on 1% agarose gels containing formaldehyde. After electrophoresis the RNA was transferred to nylon membranes (Hybond N, Amersham, UK) by the capillary transfer method (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)). Fixation of nucleic acids at the membrane was achieved by UV crosslinking (Stratalinker by Stratagene).

Membranes were prehybridized at 42°C in hybridization buffer (25% (v/v) formamide, 250 mM sodium phosphate, pH 7.2, 250 mM sodiumchloride, 1 mM EDTA 7% (w/v) SDS, 25% (w/v) polyethyleneglycol 6000, 0,25 mg/ml sheared salmon sperm DNA) for 6 h. Afterwards hybridization was performed at 42°C over night in hybridization buffer containing a radiolabelled probe in addition. The radioactive probe was prepared by using the Random Primed DNA Labelling Kit (Boehringer Mannheim, 1004760) and the approx. 4 kb KpnI/XhoI-fragment from plasmid pAlsu-pSK according to the manufacturers manual. Membranes were washed at 50°C once for 20 min in 3xSSC (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) followed by washing once for 20 min in 0.5xSSC before exposing the membrane to an x-ray-film over night.

### Claims

1. A nucleic acid molecule encoding a protein having the biological activity of an alternansucrase, selected from the group consisting of
  - (a) nucleic acid molecules encoding at least the mature form of a protein comprising the amino acid sequence indicated in Seq. ID No. 2 or the amino acid sequence which is encoded by the cDNA contained in plasmid DSM 12666;
  - (b) nucleic acid molecules comprising the nucleotide sequence indicated in Seq. ID No. 1 or the nucleotide sequence of the cDNA contained in plasmid DSM 12666 or a corresponding ribonucleotide sequence;
  - (c) nucleic acid molecules encoding a protein, the amino acid sequence of which has a homology of at least 40% to the amino acid sequence indicated in Seq. ID No 2;
  - (d) nucleic acid molecules, one strand of which hybridizes with any one of the nucleic acid molecules as defined in (a) or (b);
  - (e) nucleic acid molecules comprising a nucleotide sequence encoding a biologically active fragment of the protein which is encoded by any one of the nucleic acid molecules as defined in (a), (b), (c) or (d); and
  - (f) nucleic acid molecules, the nucleotide sequence of which deviates on account of the degeneration of the genetic code from the sequence of the nucleic acid molecules as defined in (a), (b), (c), (d) or (e).
2. An oligonucleotide or polynucleotide which specifically hybridizes with a nucleic acid molecule of claim 1.
3. A vector containing a nucleic acid molecule according to claim 1.
4. The vector according to claim 3, wherein the nucleic acid molecule is connected in sense orientation to regulatory elements ensuring the transcription and synthesis of a translatable RNA in prokaryotic or eukaryotic cells.
5. Plasmid pAlsu-pSK deposited under the accession No. DSM 12666.

6. A host cell transformed with a nucleic acid molecule of claim 1 or a vector of claim 3 or 4 or descended from such a cell.
7. The host cell according to claim 6, which is a cell of a microorganism.
8. The host cell according to claim 6, which is an *E. coli* cell.
9. A method for preparing a protein having the biological activity of an alternansucrase or a biologically active fragment thereof, wherein a host cell of any one of claims 6 to 8 is cultured under conditions permitting the synthesis of the protein, and wherein the protein is isolated from the cultured cells and/or the culture medium.
10. A protein or biologically active fragment thereof encoded by a nucleic acid molecule of claim 1 or prepared according to the method of claim 9.
11. A transgenic plant cell transformed with a nucleic acid molecule of claim 1 or a vector of claim 3 or 4, or descended from such a cell, wherein said nucleic acid molecule encoding the protein having the biological activity of an alternansucrase is under the control of regulatory elements permitting the transcription of a translatable mRNA in plant cells.
12. A plant containing the plant cells of claim 11.
13. The plant according to claim 12, which is a useful plant.
14. The plant according to claim 12 or 13, which is a sugar-storing or starch-storing plant.
15. Propagation material of a plant according to any one of claims 12 to 14, containing the plant cells of claim 11.

16. A method for preparing alternan comprising the steps of extracting and isolating the alternan from a plant according to any one of claims 12 to 14.
17. Alternan obtainable from a plant according to any one of claims 12 to 14, from propagation material according to claim 15 or by a method according to claim 16.
18. A method for preparing alternan and/or fructose, wherein
  - (a) a host cell of any one of claim 6 to 8 secretes an alternansucrase into a saccharose-containing culture medium; and
  - (b) alternan and/or fructose is/are isolated from the culture medium.
19. The method according to claim 18, wherein the host cell is immobilized.
20. A method for preparing alternan and/or fructose, wherein
  - (a) a saccharose-containing solution is contacted with a protein of claim 10 under conditions permitting the conversion of saccharose into alternan and/or fructose; and
  - (b) alternan and/or fructose is/are isolated from the solution.
21. The method according to claim 20, wherein the protein is immobilized on a carrier material.
22. A method for preparing alternan and/or fructose, wherein
  - (a) a saccharose-containing solution is contacted with a protein of claim 10 and acceptor molecules under conditions permitting the conversion of saccharose into alternan and/or fructose; and
  - (b) alternan and/or fructose is/are isolated from the solution.
23. The method according to claim 22, wherein the acceptor molecule is selected from the group consisting of maltose, isomaltose, isomaltotriose and methyl- $\alpha$ -D-glucan.

24. The method according to claim 22 or 23, wherein the protein is immobilized.
25. Cosmetic products or food products containing alternan as specified in claim 17 or prepared according to any one of claims 16 and 18 to 24.
26. A method for preparing cosmetic products or food products comprising a process according to any of claims 16 and 18 to 24 and the formulation of the resulting alternan in a form suitable for use as a cosmetic product or foodstuff.



1/12




Fig. 1

2/12




Fig. 2

3/12




Fig. 3

4/12




Fig. 4




Fig. 5

6/12




Fig. 6

7/12




Fig. 7

8/12




Fig. 8



9/12




Fig. 9

10/12




Fig. 10

11/12




Fig. 11




Fig. 12

## SEQUENCE LISTING

&lt;110&gt; MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN e.V.

&lt;120&gt; Nucleic Acid Molecules Encoding Alternansucrase

&lt;130&gt; D 1042 PCT

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 54

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 9321

&lt;212&gt; DNA

&lt;213&gt; Leuconostoc mesenteroides

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (678)..(6848)

&lt;400&gt; 1

```

gatcaaataag atgctaccgt tatacgagac gttaaggatt atcccaatct agtcgttttg 60
cgtaccttgt cgaaagctta tggctctggca aacctgagaa tcggctatgg cgtcatgcag 120
gaaccacttt atcagggttat gcaggccgta cgtttaccat ataatttaaa tacctatcaa 180
atcacagggtg cagtagctgc ccttagtgat caactttatc tgcaatcagt tgttgctaag 240
gtgaagtctg aacgtgaaaa atttgaacaa tttttgacga aacaccagtt taagtattat 300
caatcacaaa ccaactttct ctggattaaa gttggtgatg cgaaacgtgt tggtgaggct 360
cttctgtcag aagggtatca aattaatgac cgcctaaatg ccgaatggat tcgcattgca 420
ttaggaactg tgtctgataa tgaggggatg cagcgcattt tattgaattg ttaaaaaata 480
gctaagagag tatgttcttc tcttacctat ttttatttgt aattcctatt atttaatttt 540
gcatgacaat attaatagcg tgttacgatt ctactattta atgttaataa aattaataaa 600
tatggtatta tcttatatgg gtgatagatg caccaaatac tgtatcatgt ctggtcacat 660
gaaagggaga ataatta atg aaa caa caa gaa aca gtt acc cgt aaa aaa 710
                Met Lys Gln Gln Glu Thr Val Thr Arg Lys Lys
                  1                      5                      10

ctt tat aaa tcc ggt aag gtt tgg gtt gca gca gct act gca ttt gcg 758
Leu Tyr Lys Ser Gly Lys Val Trp Val Ala Ala Ala Thr Ala Phe Ala
                15                      20                      25

gta ttg ggg gtt tca act gta aca aca gtc cat gcg gat aca aat tcg 806
Val Leu Gly Val Ser Thr Val Thr Val His Ala Asp Thr Asn Ser
                30                      35                      40

aat gtc gct gtt aag caa ata aat aat aca gga acc aat gat tct ggc 854

```

2/36

Asn	Val	Ala	Val	Lys	Gln	Ile	Asn	Asn	Thr	Gly	Thr	Asn	Asp	Ser	Gly		
	45					50					55						
gaa	aaa	aag	gta	ccg	ggt	cca	tca	act	aat	aat	gat	agt	ttg	aag	caa	902	
Glu	Lys	Lys	Val	Pro	Val	Pro	Ser	Thr	Asn	Asn	Asp	Ser	Leu	Lys	Gln		
60					65					70					75		
gga	aca	gat	ggt	ttt	tgg	tat	gat	tca	gac	ggc	aat	cgt	gtc	gat	cag	950	
Gly	Thr	Asp	Gly	Phe	Trp	Tyr	Asp	Ser	Asp	Gly	Asn	Arg	Val	Asp	Gln		
				80					85					90			
aag	acc	aat	cag	att	ctg	ctt	act	gcg	gaa	caa	ctt	aaa	aaa	aat	aac	998	
Lys	Thr	Asn	Gln	Ile	Leu	Leu	Thr	Ala	Glu	Gln	Leu	Lys	Lys	Asn	Asn		
			95					100					105				
gaa	aaa	aat	tta	tca	gta	atc	agt	gat	gat	aca	tca	aaa	aaa	gat	gat	1046	
Glu	Lys	Asn	Leu	Ser	Val	Ile	Ser	Asp	Asp	Thr	Ser	Lys	Lys	Asp	Asp		
	110						115					120					
gaa	aat	att	tct	aag	cag	acc	aaa	att	gct	aat	caa	caa	aca	gta	gat	1094	
Glu	Asn	Ile	Ser	Lys	Gln	Thr	Lys	Ile	Ala	Asn	Gln	Gln	Thr	Val	Asp		
	125					130					135						
act	gct	aaa	ggc	ctg	act	acc	agt	aat	tta	tct	gat	ccc	atc	act	ggg	1142	
Thr	Ala	Lys	Gly	Leu	Thr	Thr	Ser	Asn	Leu	Ser	Asp	Pro	Ile	Thr	Gly		
140					145					150					155		
ggt	cac	tat	gaa	aat	cac	aat	ggc	tac	ttt	gtt	tat	ata	gat	gct	tca	1190	
Gly	His	Tyr	Glu	Asn	His	Asn	Gly	Tyr	Phe	Val	Tyr	Ile	Asp	Ala	Ser		
				160				165						170			
gga	aaa	caa	gta	aca	ggt	ttg	caa	aat	att	gat	ggt	aat	tta	caa	tat	1238	
Gly	Lys	Gln	Val	Thr	Gly	Leu	Gln	Asn	Ile	Asp	Gly	Asn	Leu	Gln	Tyr		
			175					180					185				
ttt	gat	gac	aat	gga	tat	caa	gtc	aag	gga	tcc	ttc	cga	gat	gtc	aac	1286	
Phe	Asp	Asp	Asn	Gly	Tyr	Gln	Val	Lys	Gly	Ser	Phe	Arg	Asp	Val	Asn		
	190						195					200					
ggc	aag	cat	atc	tat	ttt	gat	tca	gta	aca	ggg	aaa	gct	agt	tca	aat	1334	
Gly	Lys	His	Ile	Tyr	Phe	Asp	Ser	Val	Thr	Gly	Lys	Ala	Ser	Ser	Asn		
	205					210					215						
gtt	gat	att	gtt	aac	ggt	aaa	gct	caa	gga	tat	gat	gcg	caa	ggc	aac	1382	
Val	Asp	Ile	Val	Asn	Gly	Lys	Ala	Gln	Gly	Tyr	Asp	Ala	Gln	Gly	Asn		
220					225					230					235		
caa	tta	aag	aaa	agt	tat	gtc	gcc	gat	agt	tct	ggg	caa	act	tac	tat	1430	
Gln	Leu	Lys	Lys	Ser	Tyr	Val	Ala	Asp	Ser	Ser	Gly	Gln	Thr	Tyr	Tyr		
				240				245						250			
ttt	gat	ggt	aat	ggc	caa	ccg	tta	atc	ggc	ttg	caa	aca	att	gat	ggg	1478	
Phe	Asp	Gly	Asn	Gly	Gln	Pro	Leu	Ile	Gly	Leu	Gln	Thr	Ile	Asp	Gly		
			255					260					265				
aac	cta	caa	tat	ttt	aac	caa	caa	ggg	gtt	caa	ata	aag	ggt	ggt	ttc	1526	
Asn	Leu	Gln	Tyr	Phe	Asn	Gln	Gln	Gly	Val	Gln	Ile	Lys	Gly	Gly	Phe		
		270					275					280					
caa	gat	gtt	aac	aat	aaa	cgt	att	tat	ttt	gca	cca	aac	aca	ggt	aat	1574	

3/36

Gln	Asp	Val	Asn	Asn	Lys	Arg	Ile	Tyr	Phe	Ala	Pro	Asn	Thr	Gly	Asn		
285						290					295						
gcc	gtt	gcc	aat	act	gaa	ata	att	aac	ggg	aaa	tta	cag	ggg	cgt	gac	1622	
Ala	Val	Ala	Asn	Thr	Glu	Ile	Ile	Asn	Gly	Lys	Leu	Gln	Gly	Arg	Asp		
300					305				310						315		
gca	aat	ggg	aac	cag	gta	aag	aat	gca	ttt	agt	aaa	gat	gtt	gca	gga	1670	
Ala	Asn	Gly	Asn	Gln	Val	Lys	Asn	Ala	Phe	Ser	Lys	Asp	Val	Ala	Gly		
				320					325						330		
aat	aca	ttt	tat	ttt	gac	gca	aac	ggg	gtg	atg	tta	aca	ggg	ttg	caa	1718	
Asn	Thr	Phe	Tyr	Phe	Asp	Ala	Asn	Gly	Val	Met	Leu	Thr	Gly	Leu	Gln		
			335					340							345		
act	att	tca	gga	aag	aca	tat	tat	ctt	gat	gaa	caa	gga	cac	ctg	aga	1766	
Thr	Ile	Ser	Gly	Lys	Thr	Tyr	Tyr	Leu	Asp	Glu	Gln	Gly	His	Leu	Arg		
		350					355								360		
aaa	aat	tac	gcg	gga	aca	ttc	aat	aat	cag	ttt	atg	tac	ttc	gat	gct	1814	
Lys	Asn	Tyr	Ala	Gly	Thr	Phe	Asn	Asn	Gln	Phe	Met	Tyr	Phe	Asp	Ala		
		365				370					375						
gat	aca	ggg	gcg	ggg	aaa	aca	gcg	att	gaa	tat	caa	ttt	gat	caa	gga	1862	
Asp	Thr	Gly	Ala	Gly	Lys	Thr	Ala	Ile	Glu	Tyr	Gln	Phe	Asp	Gln	Gly		
380					385				390						395		
ttg	gta	tca	caa	agt	aat	gaa	aat	act	cct	cac	aat	gcc	gca	aag	tct	1910	
Leu	Val	Ser	Gln	Ser	Asn	Glu	Asn	Thr	Pro	His	Asn	Ala	Ala	Lys	Ser		
				400					405						410		
tat	gat	aaa	agt	agt	ttt	gaa	aat	gtt	gat	ggg	tac	tta	aca	gca	gat	1958	
Tyr	Asp	Lys	Ser	Ser	Phe	Glu	Asn	Val	Asp	Gly	Tyr	Leu	Thr	Ala	Asp		
			415					420							425		
aca	tgg	tat	cgt	cca	acc	gat	att	tta	aaa	aat	gga	gat	act	tgg	acg	2006	
Thr	Trp	Tyr	Arg	Pro	Thr	Asp	Ile	Leu	Lys	Asn	Gly	Asp	Thr	Trp	Thr		
		430					435								440		
gca	tct	acc	gaa	act	gat	atg	cgt	ccg	ctt	tta	atg	aca	tgg	tgg	cct	2054	
Ala	Ser	Thr	Glu	Thr	Asp	Met	Arg	Pro	Leu	Leu	Met	Thr	Trp	Trp	Pro		
		445				450											
gac	aaa	caa	aca	caa	gca	aat	tac	ttg	aat	ttt	atg	tct	agt	aaa	gga	2102	
Asp	Lys	Gln	Thr	Gln	Ala	Asn	Tyr	Leu	Asn	Phe	Met	Ser	Ser	Lys	Gly		
460					465					470					475		
ctt	ggg	ata	acg	acc	act	tat	aca	gca	gct	acg	tca	caa	aaa	aca	cta	2150	
Leu	Gly	Ile	Thr	Thr	Thr	Tyr	Thr	Ala	Ala	Thr	Ser	Gln	Lys	Thr	Leu		
				480					485						490		
aat	gac	gca	gcc	ttt	gtt	att	caa	aca	gca	att	gaa	caa	caa	ata	tct	2198	
Asn	Asp	Ala	Ala	Phe	Val	Ile	Gln	Thr	Ala	Ile	Glu	Gln	Gln	Ile	Ser		
			495					500							505		
ttg	aaa	aaa	agt	act	gag	tgg	tta	cgt	gat	gca	att	gat	agt	ttt	gtg	2246	
Leu	Lys	Lys	Ser	Thr	Glu	Trp	Leu	Arg	Asp	Ala	Ile	Asp	Ser	Phe	Val		
		510					515					520					
aag	acg	caa	gct	aat	tgg	aat	aag	caa	aca	gaa	gat	gaa	gct	ttc	gat	2294	

4/36

Lys	Thr	Gln	Ala	Asn	Trp	Asn	Lys	Gln	Thr	Glu	Asp	Glu	Ala	Phe	Asp		
525						530					535						
ggt	ttg	cag	tgg	ctt	caa	ggg	gga	ttc	cta	gct	tat	caa	gat	gat	tca	2342	
Gly	Leu	Gln	Trp	Leu	Gln	Gly	Gly	Phe	Leu	Ala	Tyr	Gln	Asp	Asp	Ser		
540					545					550					555		
cat	cgg	acg	ccg	aat	act	gat	tca	gga	aat	aac	aga	aaa	cta	gga	cgt	2390	
His	Arg	Thr	Pro	Asn	Thr	Asp	Ser	Gly	Asn	Asn	Arg	Lys	Leu	Gly	Arg		
				560					565					570			
caa	cca	att	aat	atc	gat	ggt	tcg	aaa	gat	aca	act	gat	ggt	aaa	ggc	2438	
Gln	Pro	Ile	Asn	Ile	Asp	Gly	Ser	Lys	Asp	Thr	Thr	Asp	Gly	Lys	Gly		
			575					580					585				
tct	gaa	ttc	tta	tta	gct	aac	gat	att	gac	aac	tca	aat	ccg	att	gtt	2486	
Ser	Glu	Phe	Leu	Leu	Ala	Asn	Asp	Ile	Asp	Asn	Ser	Asn	Pro	Ile	Val		
	590						595					600					
caa	gct	gag	caa	tta	aac	tgg	cta	cac	tat	tta	atg	aat	ttt	ggt	agt	2534	
Gln	Ala	Glu	Gln	Leu	Asn	Trp	Leu	His	Tyr	Leu	Met	Asn	Phe	Gly	Ser		
605						610					615						
att	aca	ggt	aat	aat	gac	aat	gcg	aat	ttt	gat	ggc	att	cgt	gta	gat	2582	
Ile	Thr	Gly	Asn	Asn	Asp	Asn	Ala	Asn	Phe	Asp	Gly	Ile	Arg	Val	Asp		
620					625					630					635		
gct	gtt	gat	aat	gtt	gat	gct	gat	tta	cta	aaa	ata	gct	ggc	gat	tat	2630	
Ala	Val	Asp	Asn	Val	Asp	Ala	Asp	Leu	Leu	Lys	Ile	Ala	Gly	Asp	Tyr		
				640					645					650			
ttt	aaa	gct	cta	tat	ggt	aca	gat	aaa	agc	gac	gcc	aat	gcc	aat	aag	2678	
Phe	Lys	Ala	Leu	Tyr	Gly	Thr	Asp	Lys	Ser	Asp	Ala	Asn	Ala	Asn	Lys		
			655					660					665				
cat	ttg	tct	att	tta	gaa	gac	tgg	aac	ggt	aaa	gat	cct	cag	tat	gtt	2726	
His	Leu	Ser	Ile	Leu	Glu	Asp	Trp	Asn	Gly	Lys	Asp	Pro	Gln	Tyr	Val		
		670					675					680					
aat	caa	cag	ggc	aat	gcg	caa	tta	aca	atg	gat	tac	aca	gtt	act	tca	2774	
Asn	Gln	Gln	Gly	Asn	Ala	Gln	Leu	Thr	Met	Asp	Tyr	Thr	Val	Thr	Ser		
685						690					695						
cag	ttt	ggc	aat	tct	cta	aca	cat	ggc	gcc	aac	aac	agg	agt	aac	atg	2822	
Gln	Phe	Gly	Asn	Ser	Leu	Thr	His	Gly	Ala	Asn	Asn	Arg	Ser	Asn	Met		
700					705				710						715		
tgg	tat	ttc	tta	gat	act	ggc	tat	tat	ctt	aat	gga	gat	ctt	aat	aag	2870	
Trp	Tyr	Phe	Leu	Asp	Thr	Gly	Tyr	Tyr	Leu	Asn	Gly	Asp	Leu	Asn	Lys		
				720				725					730				
aag	ata	gta	gat	aag	aac	cgt	cca	aat	tct	ggc	act	ttg	gtt	aac	aga	2918	
Lys	Ile	Val	Asp	Lys	Asn	Arg	Pro	Asn	Ser	Gly	Thr	Leu	Val	Asn	Arg		
			735					740					745				
att	gct	aat	tca	ggt	gat	aca	aaa	gtt	att	cca	aat	tat	agt	ttt	gtt	2966	
Ile	Ala	Asn	Ser	Gly	Asp	Thr	Lys	Val	Ile	Pro	Asn	Tyr	Ser	Phe	Val		
		750					755					760					
aga	gca	cat	gat	tac	gat	gct	caa	gat	cca	att	aga	aaa	gcc	atg	att	3014	



5/36

Arg	Ala	His	Asp	Tyr	Asp	Ala	Gln	Asp	Pro	Ile	Arg	Lys	Ala	Met	Ile		
765						770					775						
gat	cat	ggg	att	att	aaa	aac	atg	cag	gat	act	ttc	act	ttt	gac	caa	3062	
Asp	His	Gly	Ile	Ile	Lys	Asn	Met	Gln	Asp	Thr	Phe	Thr	Phe	Asp	Gln		
780					785					790					795		
ctg	gct	cag	gga	atg	gaa	ttc	tac	tat	aaa	gat	caa	gag	aat	ccg	tct	3110	
Leu	Ala	Gln	Gly	Met	Glu	Phe	Tyr	Tyr	Lys	Asp	Gln	Glu	Asn	Pro	Ser		
				800					805						810		
ggg	ttc	aaa	aag	tat	aac	gat	tat	aac	tta	cct	agt	gct	tat	gca	atg	3158	
Gly	Phe	Lys	Lys	Tyr	Asn	Asp	Tyr	Asn	Leu	Pro	Ser	Ala	Tyr	Ala	Met		
			815					820						825			
ttg	ttg	act	aat	aag	gat	act	gta	cct	cgt	gtc	tat	tat	gga	gat	atg	3206	
Leu	Leu	Thr	Asn	Lys	Asp	Thr	Val	Pro	Arg	Val	Tyr	Tyr	Gly	Asp	Met		
			830				835							840			
tac	ctc	gaa	ggc	ggg	caa	tat	atg	gaa	aaa	ggg	acg	att	tac	aat	cct	3254	
Tyr	Leu	Glu	Gly	Gly	Gln	Tyr	Met	Glu	Lys	Gly	Thr	Ile	Tyr	Asn	Pro		
	845					850					855						
gtc	att	tca	gcg	ttg	ctc	aaa	gct	aga	ata	aaa	tat	gtt	tct	ggg	ggg	3302	
Val	Ile	Ser	Ala	Leu	Leu	Lys	Ala	Arg	Ile	Lys	Tyr	Val	Ser	Gly	Gly		
860					865					870					875		
caa	aca	atg	gct	acc	gat	agt	tct	gga	aaa	gac	ctt	aaa	gat	ggc	gaa	3350	
Gln	Thr	Met	Ala	Thr	Asp	Ser	Ser	Gly	Lys	Asp	Leu	Lys	Asp	Gly	Glu		
				880					885						890		
act	gat	ttg	tta	aca	agt	gtt	cga	ttt	ggg	aaa	gga	att	atg	aca	tca	3398	
Thr	Asp	Leu	Leu	Thr	Ser	Val	Arg	Phe	Gly	Lys	Gly	Ile	Met	Thr	Ser		
				895				900						905			
gat	caa	acc	aca	aca	caa	gac	aat	agc	caa	gat	tat	aaa	aat	caa	ggc	3446	
Asp	Gln	Thr	Thr	Thr	Gln	Asp	Asn	Ser	Gln	Asp	Tyr	Lys	Asn	Gln	Gly		
		910					915						920				
atc	ggg	gtc	att	gtt	ggg	aat	aac	cct	gac	ctt	aag	ttg	aac	aat	gat	3494	
Ile	Gly	Val	Ile	Val	Gly	Asn	Asn	Pro	Asp	Leu	Lys	Leu	Asn	Asn	Asp		
	925					930					935						
aag	acc	att	acc	ttg	cat	atg	gga	aag	gcg	cat	aag	aat	caa	ctt	tac	3542	
Lys	Thr	Ile	Thr	Leu	His	Met	Gly	Lys	Ala	His	Lys	Asn	Gln	Leu	Tyr		
940					945					950					955		
cgt	gcc	tta	gta	tta	tca	aat	gac	tca	gga	att	gat	gtt	tat	gat	agt	3590	
Arg	Ala	Leu	Val	Leu	Ser	Asn	Asp	Ser	Gly	Ile	Asp	Val	Tyr	Asp	Ser		
				960					965						970		
gat	gat	aaa	gca	cca	act	ttg	aga	aca	aat	gac	aac	ggg	gac	ttg	att	3638	
Asp	Asp	Lys	Ala	Pro	Thr	Leu	Arg	Thr	Asn	Asp	Asn	Gly	Asp	Leu	Ile		
			975					980						985			
ttc	cat	aag	aca	aat	acg	ttt	gtg	aag	caa	gat	gga	act	att	ata	aat	3686	
Phe	His	Lys	Thr	Asn	Thr	Phe	Val	Lys	Gln	Asp	Gly	Thr	Ile	Ile	Asn		
		990					995					1000					
tac	gaa	atg	aag	gga	tca	tta	aat	gct	tta	att	tca	ggg	tat	tta	ggg	3734	

6/36

Tyr Glu Met Lys Gly Ser Leu Asn Ala Leu Ile Ser Gly Tyr Leu Gly	
1005	1010 1015
gtc tgg gtg cca gtt gga gct agt gat tca caa gat gct cgt aca gtg	3782
Val Trp Val Pro Val Gly Ala Ser Asp Ser Gln Asp Ala Arg Thr Val	
1020	1025 1030 1035
gca act gag tca tca tca agt aat gat ggt tct gta ttc cat tca aat	3830
Ala Thr Glu Ser Ser Ser Ser Asn Asp Gly Ser Val Phe His Ser Asn	
	1040 1045 1050
gct gca tta gat tct aat gtt ata tat gaa ggc ttt tca aac ttt caa	3878
Ala Ala Leu Asp Ser Asn Val Ile Tyr Glu Gly Phe Ser Asn Phe Gln	
	1055 1060 1065
gcg atg ccg act tct cct gag caa agt aca aat gtt gtt att gca aca	3926
Ala Met Pro Thr Ser Pro Glu Gln Ser Thr Asn Val Val Ile Ala Thr	
	1070 1075 1080
aag gct aac tta ttt aaa gaa tta ggt att act agt ttt gag tta gca	3974
Lys Ala Asn Leu Phe Lys Glu Leu Gly Ile Thr Ser Phe Glu Leu Ala	
	1085 1090 1095
cct caa tat agg tct agt ggt gac act aat tac ggt ggc atg tca ttc	4022
Pro Gln Tyr Arg Ser Ser Gly Asp Thr Asn Tyr Gly Gly Met Ser Phe	
1100	1105 1110 1115
tta gat tct ttc tta aat aat ggt tat gca ttt acc gat aga tat gat	4070
Leu Asp Ser Phe Leu Asn Asn Gly Tyr Ala Phe Thr Asp Arg Tyr Asp	
	1120 1125 1130
tta ggc ttt aac aaa gca gac ggg aat cct aac cca aca aag tat gga	4118
Leu Gly Phe Asn Lys Ala Asp Gly Asn Pro Asn Pro Thr Lys Tyr Gly	
	1135 1140 1145
aca gat caa gat tta cgt aat gca ata gag gca tta cac aaa aac ggc	4166
Thr Asp Gln Asp Leu Arg Asn Ala Ile Glu Ala Leu His Lys Asn Gly	
	1150 1155 1160
atg cag gct ata gct gat tgg gtt cct gac caa ata tat gct tta cca	4214
Met Gln Ala Ile Ala Asp Trp Val Pro Asp Gln Ile Tyr Ala Leu Pro	
	1165 1170 1175
gga aag gaa gtt gtt acc gct act aga gta gac gaa cgg gga aat caa	4262
Gly Lys Glu Val Val Thr Ala Thr Arg Val Asp Glu Arg Gly Asn Gln	
1180	1185 1190 1195
cta aaa gac aca gat ttt gtc aac tta ctc tat gtt gct aat act aaa	4310
Leu Lys Asp Thr Asp Phe Val Asn Leu Leu Tyr Val Ala Asn Thr Lys	
	1200 1205 1210
agt agt ggt gtg gat tat cag gca aag tat ggc ggc gaa ttt tta gat	4358
Ser Ser Gly Val Asp Tyr Gln Ala Lys Tyr Gly Gly Glu Phe Leu Asp	
	1215 1220 1225
aaa tta aga gaa gag tac cca tcg tta ttc aaa cag aac caa gta tcg	4406
Lys Leu Arg Glu Glu Tyr Pro Ser Leu Phe Lys Gln Asn Gln Val Ser	
	1230 1235 1240
aca ggt cag cca att gat gct tct aca aaa att aag caa tgg tca gct	4454

7/36

Thr Gly Gln Pro Ile Asp Ala Ser Thr Lys Ile Lys Gln Trp Ser Ala	
1245 1250 1255	
aaa tat atg aat ggg acc aat att tta cat cga ggt gct tat tat gtt	4502
Lys Tyr Met Asn Gly Thr Asn Ile Leu His Arg Gly Ala Tyr Tyr Val	
1260 1265 1270 1275	
ttg aaa gac tgg gct act aac cag tat ttt aac att gca aaa acg aat	4550
Leu Lys Asp Trp Ala Thr Asn Gln Tyr Phe Asn Ile Ala Lys Thr Asn	
1280 1285 1290	
gaa gta ttt ttg cca cta cag ttg cag aat aaa gat gcg caa act ggt	4598
Glu Val Phe Leu Pro Leu Gln Leu Gln Asn Lys Asp Ala Gln Thr Gly	
1295 1300 1305	
ttc att agt gat gcc tcc ggt gta aaa tat tac tca att agt ggt tat	4646
Phe Ile Ser Asp Ala Ser Gly Val Lys Tyr Tyr Ser Ile Ser Gly Tyr	
1310 1315 1320	
caa gca aaa gat act ttt att gaa gat ggt aat ggg aat tgg tat tac	4694
Gln Ala Lys Asp Thr Phe Ile Glu Asp Gly Asn Gly Asn Trp Tyr Tyr	
1325 1330 1335	
ttt gat aaa gat ggt tac atg gtg cgt tgc cag caa gga gaa aat cct	4742
Phe Asp Lys Asp Gly Tyr Met Val Arg Ser Gln Gln Gly Glu Asn Pro	
1340 1345 1350 1355	
ata aga aca gtc gaa act agt gtc aac aca cga aac ggt aat tat tac	4790
Ile Arg Thr Val Glu Thr Ser Val Asn Thr Arg Asn Gly Asn Tyr Tyr	
1360 1365 1370	
ttt atg cca aat ggt gtc gag ttg cgc aaa ggc ttt gga acg gat aat	4838
Phe Met Pro Asn Gly Val Glu Leu Arg Lys Gly Phe Gly Thr Asp Asn	
1375 1380 1385	
agt ggt aat gtc tat tat ttt gat gat caa ggt aag atg gtg aga gat	4886
Ser Gly Asn Val Tyr Tyr Phe Asp Asp Gln Gly Lys Met Val Arg Asp	
1390 1395 1400	
aaa tac att aac gat gat gct aat aat ttt tat cac tta aat gtt gat	4934
Lys Tyr Ile Asn Asp Asp Ala Asn Asn Phe Tyr His Leu Asn Val Asp	
1405 1410 1415	
ggg act atg tct cga gga cta ttt aaa ttt gat tct gat act cta cag	4982
Gly Thr Met Ser Arg Gly Leu Phe Lys Phe Asp Ser Asp Thr Leu Gln	
1420 1425 1430 1435	
tat ttt gct agt aat ggt gtc caa ata aaa gat agt tat gcg aag gat	5030
Tyr Phe Ala Ser Asn Gly Val Gln Ile Lys Asp Ser Tyr Ala Lys Asp	
1440 1445 1450	
agt aaa ggc aat aaa tat tat ttt gac tca gct aca gga aat aac gat	5078
Ser Lys Gly Asn Lys Tyr Tyr Phe Asp Ser Ala Thr Gly Asn Asn Asp	
1455 1460 1465	
act ggg aaa gcc caa act tgg gat ggt aat ggc tac tat att act att	5126
Thr Gly Lys Ala Gln Thr Trp Asp Gly Asn Gly Tyr Tyr Ile Thr Ile	
1470 1475 1480	
gat tct gat gcg aac aat aca att ggg gtt aac aca gac tac act gcc	5174

Asp Ser Asp Ala Asn Asn Thr Ile Gly Val Asn Thr Asp Tyr Thr Ala	
1485 1490 1495	
tac atc act agc tcg ctg cgc gaa gat ggc tta ttt gct aac gca cct	5222
Tyr Ile Thr Ser Ser Leu Arg Glu Asp Gly Leu Phe Ala Asn Ala Pro	
1500 1505 1510 1515	
tac ggt gtt gta aca aaa gac caa aat ggt aac gat ctt aag tgg cag	5270
Tyr Gly Val Val Thr Lys Asp Gln Asn Gly Asn Asp Leu Lys Trp Gln	
1520 1525 1530	
tat att aac cat acg aaa cag tac gaa ggg caa caa gtg caa gtc acg	5318
Tyr Ile Asn His Thr Lys Gln Tyr Glu Gly Gln Gln Val Gln Val Thr	
1535 1540 1545	
cgt caa tac aca gac agt aag gga gtc agc tgg aac tta att acc ttt	5366
Arg Gln Tyr Thr Asp Ser Lys Gly Val Ser Trp Asn Leu Ile Thr Phe	
1550 1555 1560	
gct ggt ggt gat tta caa gga caa agg ctt tgg gtg gat agt cgt gcg	5414
Ala Gly Gly Asp Leu Gln Gly Gln Arg Leu Trp Val Asp Ser Arg Ala	
1565 1570 1575	
tta act atg aca cca ttt aaa acg atg aac caa ata agc ttc att agt	5462
Leu Thr Met Thr Pro Phe Lys Thr Met Asn Gln Ile Ser Phe Ile Ser	
1580 1585 1590 1595	
tat gct aac cgc aat gat ggg ttg ttt ttg aat gcg cca tac caa gtc	5510
Tyr Ala Asn Arg Asn Asp Gly Leu Phe Leu Asn Ala Pro Tyr Gln Val	
1600 1605 1610	
aag ggg tat caa tta gct ggg atg tcc aac caa tac aag ggc caa caa	5558
Lys Gly Tyr Gln Leu Ala Gly Met Ser Asn Gln Tyr Lys Gly Gln Gln	
1615 1620 1625	
gtg acc att gct ggg gtg gcg aac gtt tct gga aaa gac tgg agt ctg	5606
Val Thr Ile Ala Gly Val Ala Asn Val Ser Gly Lys Asp Trp Ser Leu	
1630 1635 1640	
att agt ttt aat ggg aca cag tac tgg att gat agt cag gca ttg aat	5654
Ile Ser Phe Asn Gly Thr Gln Tyr Trp Ile Asp Ser Gln Ala Leu Asn	
1645 1650 1655	
acc aat ttc aca cat gac atg aac caa aag gtc ttt gtc aat aca act	5702
Thr Asn Phe Thr His Asp Met Asn Gln Lys Val Phe Val Asn Thr Thr	
1660 1665 1670 1675	
agt aat ctt gat ggg tta ttc tta aat gcg cca tac cgt caa ccg ggt	5750
Ser Asn Leu Asp Gly Leu Phe Leu Asn Ala Pro Tyr Arg Gln Pro Gly	
1680 1685 1690	
tat aag tta gcc ggt ttg gct aaa aat tac aac aac caa acg gtt act	5798
Tyr Lys Leu Ala Gly Leu Ala Lys Asn Tyr Asn Asn Gln Thr Val Thr	
1695 1700 1705	
gtt agt caa cag tac ttt gat gat caa ggc acg gtc tgg agt cag gtt	5846
Val Ser Gln Gln Tyr Phe Asp Asp Gln Gly Thr Val Trp Ser Gln Val	
1710 1715 1720	
gtc ctt ggg ggt cag acg gtc tgg gtt gat aac cat gca ttg gca cag	5894

9/36

Val Leu Gly Gly Gln Thr Val Trp Val Asp Asn His Ala Leu Ala Gln	
1725 1730 1735	
atg caa gtt agt gat aca gac caa cag ctc tat gtg aat agc aat ggt	5942
Met Gln Val Ser Asp Thr Asp Gln Gln Leu Tyr Val Asn Ser Asn Gly	
1740 1745 1750 1755	
cgg aat gat ggg tta ttc ttg aat gcg cca tat cgt ggt caa ggg tca	5990
Arg Asn Asp Gly Leu Phe Leu Asn Ala Pro Tyr Arg Gly Gln Gly Ser	
1760 1765 1770	
caa ctg ata ggc atg acg gca gat tat aat ggg caa cat gta caa gtg	6038
Gln Leu Ile Gly Met Thr Ala Asp Tyr Asn Gly Gln His Val Gln Val	
1775 1780 1785	
acc aag caa ggg caa gat gcc tat ggt gca caa tgg cgt ctt att acg	6086
Thr Lys Gln Gly Gln Asp Ala Tyr Gly Ala Gln Trp Arg Leu Ile Thr	
1790 1795 1800	
cta aat aat caa cag gtc tgg gtt gat agt cgc gct ttg agc aca aca	6134
Leu Asn Asn Gln Gln Val Trp Val Asp Ser Arg Ala Leu Ser Thr Thr	
1805 1810 1815	
atc atg caa gcc atg aat gat aat atg tat gta aat agc agc caa cgg	6182
Ile Met Gln Ala Met Asn Asp Asn Met Tyr Val Asn Ser Ser Gln Arg	
1820 1825 1830 1835	
aca gat ggc ttg tgg tta aac gca cct tat acg atg agt ggg gct aaa	6230
Thr Asp Gly Leu Trp Leu Asn Ala Pro Tyr Thr Met Ser Gly Ala Lys	
1840 1845 1850	
tgg gct ggt gat aca cgt tca gct aat ggg cgc tat gtc cat att tca	6278
Trp Ala Gly Asp Thr Arg Ser Ala Asn Gly Arg Tyr Val His Ile Ser	
1855 1860 1865	
aaa gct tat tca aac gaa gtc ggc aat aca tat tac ttg acg aat ttg	6326
Lys Ala Tyr Ser Asn Glu Val Gly Asn Thr Tyr Tyr Leu Thr Asn Leu	
1870 1875 1880	
aat ggt caa agc aca tgg att gac aag cgg gcg ttt act gtg acc ttc	6374
Asn Gly Gln Ser Thr Trp Ile Asp Lys Arg Ala Phe Thr Val Thr Phe	
1885 1890 1895	
gat cag gtg gtg gca tta aat gca acg att gtg gca cgc caa cga cca	6422
Asp Gln Val Val Ala Leu Asn Ala Thr Ile Val Ala Arg Gln Arg Pro	
1900 1905 1910 1915	
gat ggg atg ttt aag aca gca cca tat ggt gaa gcg ggg gcg cag ttt	6470
Asp Gly Met Phe Lys Thr Ala Pro Tyr Gly Glu Ala Gly Ala Gln Phe	
1920 1925 1930	
gtc gat tat gtg aca aac tat aac cag caa acc gtg cca gta aca aag	6518
Val Asp Tyr Val Thr Asn Tyr Asn Gln Gln Thr Val Pro Val Thr Lys	
1935 1940 1945	
caa cat tca gat gct cag ggg aat caa tgg tac tta gcg aca gtg aat	6566
Gln His Ser Asp Ala Gln Gly Asn Gln Trp Tyr Leu Ala Thr Val Asn	
1950 1955 1960	
ggg aca caa tac tgg att gat caa cgg tca ttt tca cca gta gta acg	6614

10/36

Gly Thr Gln Tyr Trp Ile Asp Gln Arg Ser Phe Ser Pro Val Val Thr	
1965 1970 1975	
aag gtg gtt gat tat caa gct aag att gtg cca cgg aca aca cgt gat	6662
Lys Val Val Asp Tyr Gln Ala Lys Ile Val Pro Arg Thr Thr Arg Asp	
1980 1985 1990 1995	
ggt gtg ttt agt ggc gca ccc tat ggg gaa gtg aat gct aag cta gtt	6710
Gly Val Phe Ser Gly Ala Pro Tyr Gly Glu Val Asn Ala Lys Leu Val	
2000 2005 2010	
aac atg gca act gcg tat caa aat caa gtt gtc cat gcg aca ggg gaa	6758
Asn Met Ala Thr Ala Tyr Gln Asn Gln Val Val His Ala Thr Gly Glu	
2015 2020 2025	
tat acg aat gct tca ggg atc aca tgg agt cag ttc gcg tta agc ggg	6806
Tyr Thr Asn Ala Ser Gly Ile Thr Trp Ser Gln Phe Ala Leu Ser Gly	
2030 2035 2040	
caa gaa gac aag cta tgg att gat aag cgt gct ttg caa gct	6848
Gln Glu Asp Lys Leu Trp Ile Asp Lys Arg Ala Leu Gln Ala	
2045 2050 2055	
taagggaagg attcgacaaa ggagggtaac attatcagcg gatggtgtta tcctcctttc	6908
ctgtactcag tatttcccaa ataattgaga cagtttcatg acaaatcaac aaaactagtg	6968
tcaatgcctc ggttatgggg taaactacta ttagttaaag ggttggtgca taataatata	7028
acattcaata tattatgtat ttttatctga ttatgtgatt ttttgagatt tggagcgaaa	7088
atgaataaag taataattaa taaaagtttc tgtgttttgg taatgagttt gttgtcgata	7148
tttctattct cattaagagt agacgctcgt tccaatagt gctacaatca aatatctttt	7208
aatcaacatg aaattgctta tgcgcctttt agtcaagtac cttggtatgc aactgtgaat	7268
gtcggaatgt ctgcagacaa aaataatata tatacagcta ttgatatggg gccatagtat	7328
caagggaactt ctttttttcc gtgggtggtat caacatgata attatgatta tcatgatggt	7388
tatcaacttg cgattgacaa tcaatcgctt ttcataaaaa tgggtgatac aggaaggctg	7448
tcgtttgatg ttattgagtt gtatacaggt aaaaaagttg gttatggaaa attcacagcc	7508
tataactata acaagacttc atggcagttt caaatatctt taaaggcgat aaagaaaatt	7568
ttaaataata acatatacaga aaatgcgaca attagtttgt acaatgggta tctattcaaa	7628
caagccgaaa gtatcacgta tgctgggtca ccaacagggc catgggttact tgccggtatt	7688
ggtttggttt ttgctgggtg gggctattac ctgcaaaatc agcgccgaaa aacgattacg	7748
gcatcatcaa caaacgaggt aacctatgct taataatgct gttattattg gtctaattat	7808
tatttgggtc tatgtattat ctgttctgaa gcgtgctaaa acggatgcgt tttatttttt	7868
ggctggcagc gcaggcttgt tcgtcattct tcttttatgg agtaaacctt atggtgtttg	7928
gttgttttcg accattctca cttggctcggc cgggggttgt ggtcatctca ctggcttatt	7988

tgatactttc tatgctcac atgttattca ggttgtggcg aatcatcata ccagcatttt 8048  
 attggttgat tatgaatgtt caggcattat tgaacaacg gcattttggg gattaattgc 8108  
 cttttatcct gtgtatgatg cacaaaagcg gttaatgttg gcgcttattg gggcattatg 8168  
 gtttttcctt gcaaatacgt tacgactatc ttttgttgca gttgtaattt attattttgg 8228  
 cgatggcgct ttttatgttg ctcatcatt caat tatcggacgc ctgttgttct attcaattgt 8288  
 aatcttgctt tattatctag tcttcataaa ggggcaactt gttaatcaaa tgctcaaaaa 8348  
 acagggataa actatgtcat tttatttaca tcaatttact tatcagttag gtttttggtt 8408  
 cacttggttg ctcatccgt ttgttgtgga aattttccca gctattattt caatctagtt 8468  
 ttgataaaaa catcaaaaaa gcatcaagt atgcaagaac cattgaaatt accaatggta 8528  
 tctattgtat tacctatata taattcagga caaacattat accagtgtat tcagtcgatt 8588  
 agtcaatcaa cctatccaaa acaattaatc caaattattg ccgtgaataa tcaaagcact 8648  
 gataacagtt ttacagtgtt taatcaggct caagctgatt tccaataact tagaatgcag 8708  
 tggatgaaca cagatcaagg taaagcgctg gcactaaatg ctgctattta taatagtatg 8768  
 gggcaatata ttattaattt ggatactgat ggttggttag aaccatgc cttaaacgg 8828  
 tttgtgcttt attttgaaaa tcaactcagaa attgatgtag caactggtac gattctgaca 8888  
 caaaaaaaaa tgattcaaaa aacgcaaagt aaatggctta aattgctaca gttaaacgaa 8948  
 tattttgagt atgcacaatc attcttatca ggccgtagta ttgaaaaccg tggtaatcgt 9008  
 ttgttcacaa tgcagggtgc attttcagcg ttttagacgtg atgtattagt tcagacattt 9068  
 atgtataatg ttgacacggt tgggtgaagat actgatatga catttcaact gcgattccgc 9128  
 ctaggcaagc gtattggttt ctgtgatgat gccatgtttt atgttgaacc aatatcaggt 9188  
 tatagtgaac tttatttaca acggcagcgt tggcagcgtg ggcaaattga agtcgcacaa 9248  
 aatttcatgc agaataagtt gagtgtccgt cagattttta ctaactttat gattagtcga 9308  
 ttaatgattg atc 9321

&lt;210&gt; 2

&lt;211&gt; 2057

&lt;212&gt; PRT

<213> *Leuconostoc mesenteroides*

&lt;400&gt; 2

Met	Lys	Gln	Gln	Glu	Thr	Val	Thr	Arg	Lys	Lys	Leu	Tyr	Lys	Ser	Gly
1					5				10					15	

Lys	Val	Trp	Val	Ala	Ala	Ala	Thr	Ala	Phe	Ala	Val	Leu	Gly	Val	Ser
			20					25					30		

12/36

Thr Val Thr Thr Val His Ala Asp Thr Asn Ser Asn Val Ala Val Lys  
 35 40 45  
 Gln Ile Asn Asn Thr Gly Thr Asn Asp Ser Gly Glu Lys Lys Val Pro  
 50 55 60  
 Val Pro Ser Thr Asn Asn Asp Ser Leu Lys Gln Gly Thr Asp Gly Phe  
 65 70 75 80  
 Trp Tyr Asp Ser Asp Gly Asn Arg Val Asp Gln Lys Thr Asn Gln Ile  
 85 90 95  
 Leu Leu Thr Ala Glu Gln Leu Lys Lys Asn Asn Glu Lys Asn Leu Ser  
 100 105 110  
 Val Ile Ser Asp Asp Thr Ser Lys Lys Asp Asp Glu Asn Ile Ser Lys  
 115 120 125  
 Gln Thr Lys Ile Ala Asn Gln Gln Thr Val Asp Thr Ala Lys Gly Leu  
 130 135 140  
 Thr Thr Ser Asn Leu Ser Asp Pro Ile Thr Gly Gly His Tyr Glu Asn  
 145 150 155 160  
 His Asn Gly Tyr Phe Val Tyr Ile Asp Ala Ser Gly Lys Gln Val Thr  
 165 170 175  
 Gly Leu Gln Asn Ile Asp Gly Asn Leu Gln Tyr Phe Asp Asp Asn Gly  
 180 185 190  
 Tyr Gln Val Lys Gly Ser Phe Arg Asp Val Asn Gly Lys His Ile Tyr  
 195 200 205  
 Phe Asp Ser Val Thr Gly Lys Ala Ser Ser Asn Val Asp Ile Val Asn  
 210 215 220  
 Gly Lys Ala Gln Gly Tyr Asp Ala Gln Gly Asn Gln Leu Lys Lys Ser  
 225 230 235 240  
 Tyr Val Ala Asp Ser Ser Gly Gln Thr Tyr Tyr Phe Asp Gly Asn Gly  
 245 250 255  
 Gln Pro Leu Ile Gly Leu Gln Thr Ile Asp Gly Asn Leu Gln Tyr Phe  
 260 265 270  
 Asn Gln Gln Gly Val Gln Ile Lys Gly Gly Phe Gln Asp Val Asn Asn  
 275 280 285  
 Lys Arg Ile Tyr Phe Ala Pro Asn Thr Gly Asn Ala Val Ala Asn Thr  
 290 295 300  
 Glu Ile Ile Asn Gly Lys Leu Gln Gly Arg Asp Ala Asn Gly Asn Gln  
 305 310 315 320  
 Val Lys Asn Ala Phe Ser Lys Asp Val Ala Gly Asn Thr Phe Tyr Phe  
 325 330 335  
 Asp Ala Asn Gly Val Met Leu Thr Gly Leu Gln Thr Ile Ser Gly Lys  
 340 345 350



13/36

Thr Tyr Tyr Leu Asp Glu Gln Gly His Leu Arg Lys Asn Tyr Ala Gly  
 355 360 365  
 Thr Phe Asn Asn Gln Phe Met Tyr Phe Asp Ala Asp Thr Gly Ala Gly  
 370 375 380  
 Lys Thr Ala Ile Glu Tyr Gln Phe Asp Gln Gly Leu Val Ser Gln Ser  
 385 390 395 400  
 Asn Glu Asn Thr Pro His Asn Ala Ala Lys Ser Tyr Asp Lys Ser Ser  
 405 410 415  
 Phe Glu Asn Val Asp Gly Tyr Leu Thr Ala Asp Thr Trp Tyr Arg Pro  
 420 425 430  
 Thr Asp Ile Leu Lys Asn Gly Asp Thr Trp Thr Ala Ser Thr Glu Thr  
 435 440 445  
 Asp Met Arg Pro Leu Leu Met Thr Trp Trp Pro Asp Lys Gln Thr Gln  
 450 455 460  
 Ala Asn Tyr Leu Asn Phe Met Ser Ser Lys Gly Leu Gly Ile Thr Thr  
 465 470 475 480  
 Thr Tyr Thr Ala Ala Thr Ser Gln Lys Thr Leu Asn Asp Ala Ala Phe  
 485 490 495  
 Val Ile Gln Thr Ala Ile Glu Gln Gln Ile Ser Leu Lys Lys Ser Thr  
 500 505 510  
 Glu Trp Leu Arg Asp Ala Ile Asp Ser Phe Val Lys Thr Gln Ala Asn  
 515 520 525  
 Trp Asn Lys Gln Thr Glu Asp Glu Ala Phe Asp Gly Leu Gln Trp Leu  
 530 535 540  
 Gln Gly Gly Phe Leu Ala Tyr Gln Asp Asp Ser His Arg Thr Pro Asn  
 545 550 555 560  
 Thr Asp Ser Gly Asn Asn Arg Lys Leu Gly Arg Gln Pro Ile Asn Ile  
 565 570 575  
 Asp Gly Ser Lys Asp Thr Thr Asp Gly Lys Gly Ser Glu Phe Leu Leu  
 580 585 590  
 Ala Asn Asp Ile Asp Asn Ser Asn Pro Ile Val Gln Ala Glu Gln Leu  
 595 600 605  
 Asn Trp Leu His Tyr Leu Met Asn Phe Gly Ser Ile Thr Gly Asn Asn  
 610 615 620  
 Asp Asn Ala Asn Phe Asp Gly Ile Arg Val Asp Ala Val Asp Asn Val  
 625 630 635 640  
 Asp Ala Asp Leu Leu Lys Ile Ala Gly Asp Tyr Phe Lys Ala Leu Tyr  
 645 650 655  
 Gly Thr Asp Lys Ser Asp Ala Asn Ala Asn Lys His Leu Ser Ile Leu  
 660 665 670

14/36

Glu Asp Trp Asn Gly Lys Asp Pro Gln Tyr Val Asn Gln Gln Gly Asn  
 675 680 685  
 Ala Gln Leu Thr Met Asp Tyr Thr Val Thr Ser Gln Phe Gly Asn Ser  
 690 695 700  
 Leu Thr His Gly Ala Asn Asn Arg Ser Asn Met Trp Tyr Phe Leu Asp  
 705 710 715 720  
 Thr Gly Tyr Tyr Leu Asn Gly Asp Leu Asn Lys Lys Ile Val Asp Lys  
 725 730 735  
 Asn Arg Pro Asn Ser Gly Thr Leu Val Asn Arg Ile Ala Asn Ser Gly  
 740 745 750  
 Asp Thr Lys Val Ile Pro Asn Tyr Ser Phe Val Arg Ala His Asp Tyr  
 755 760 765  
 Asp Ala Gln Asp Pro Ile Arg Lys Ala Met Ile Asp His Gly Ile Ile  
 770 775 780  
 Lys Asn Met Gln Asp Thr Phe Thr Phe Asp Gln Leu Ala Gln Gly Met  
 785 790 795 800  
 Glu Phe Tyr Tyr Lys Asp Gln Glu Asn Pro Ser Gly Phe Lys Lys Tyr  
 805 810 815  
 Asn Asp Tyr Asn Leu Pro Ser Ala Tyr Ala Met Leu Leu Thr Asn Lys  
 820 825 830  
 Asp Thr Val Pro Arg Val Tyr Tyr Gly Asp Met Tyr Leu Glu Gly Gly  
 835 840 845  
 Gln Tyr Met Glu Lys Gly Thr Ile Tyr Asn Pro Val Ile Ser Ala Leu  
 850 855 860  
 Leu Lys Ala Arg Ile Lys Tyr Val Ser Gly Gly Gln Thr Met Ala Thr  
 865 870 875 880  
 Asp Ser Ser Gly Lys Asp Leu Lys Asp Gly Glu Thr Asp Leu Leu Thr  
 885 890 895  
 Ser Val Arg Phe Gly Lys Gly Ile Met Thr Ser Asp Gln Thr Thr Thr  
 900 905 910  
 Gln Asp Asn Ser Gln Asp Tyr Lys Asn Gln Gly Ile Gly Val Ile Val  
 915 920 925  
 Gly Asn Asn Pro Asp Leu Lys Leu Asn Asn Asp Lys Thr Ile Thr Leu  
 930 935 940  
 His Met Gly Lys Ala His Lys Asn Gln Leu Tyr Arg Ala Leu Val Leu  
 945 950 955 960  
 Ser Asn Asp Ser Gly Ile Asp Val Tyr Asp Ser Asp Asp Lys Ala Pro  
 965 970 975  
 Thr Leu Arg Thr Asn Asp Asn Gly Asp Leu Ile Phe His Lys Thr Asn  
 980 985 990

15/36

Thr Phe Val Lys Gln Asp Gly Thr Ile Ile Asn Tyr Glu Met Lys Gly  
 995 1000 1005  
 Ser Leu Asn Ala Leu Ile Ser Gly Tyr Leu Gly Val Trp Val Pro Val  
 1010 1015 1020  
 Gly Ala Ser Asp Ser Gln Asp Ala Arg Thr Val Ala Thr Glu Ser Ser  
 025 1030 1035 1040  
 Ser Ser Asn Asp Gly Ser Val Phe His Ser Asn Ala Ala Leu Asp Ser  
 1045 1050 1055  
 Asn Val Ile Tyr Glu Gly Phe Ser Asn Phe Gln Ala Met Pro Thr Ser  
 1060 1065 1070  
 Pro Glu Gln Ser Thr Asn Val Val Ile Ala Thr Lys Ala Asn Leu Phe  
 1075 1080 1085  
 Lys Glu Leu Gly Ile Thr Ser Phe Glu Leu Ala Pro Gln Tyr Arg Ser  
 1090 1095 1100  
 Ser Gly Asp Thr Asn Tyr Gly Gly Met Ser Phe Leu Asp Ser Phe Leu  
 105 1110 1115 1120  
 Asn Asn Gly Tyr Ala Phe Thr Asp Arg Tyr Asp Leu Gly Phe Asn Lys  
 1125 1130 1135  
 Ala Asp Gly Asn Pro Asn Pro Thr Lys Tyr Gly Thr Asp Gln Asp Leu  
 1140 1145 1150  
 Arg Asn Ala Ile Glu Ala Leu His Lys Asn Gly Met Gln Ala Ile Ala  
 1155 1160 1165  
 Asp Trp Val Pro Asp Gln Ile Tyr Ala Leu Pro Gly Lys Glu Val Val  
 1170 1175 1180  
 Thr Ala Thr Arg Val Asp Glu Arg Gly Asn Gln Leu Lys Asp Thr Asp  
 185 1190 1195 1200  
 Phe Val Asn Leu Leu Tyr Val Ala Asn Thr Lys Ser Ser Gly Val Asp  
 1205 1210 1215  
 Tyr Gln Ala Lys Tyr Gly Gly Glu Phe Leu Asp Lys Leu Arg Glu Glu  
 1220 1225 1230  
 Tyr Pro Ser Leu Phe Lys Gln Asn Gln Val Ser Thr Gly Gln Pro Ile  
 1235 1240 1245  
 Asp Ala Ser Thr Lys Ile Lys Gln Trp Ser Ala Lys Tyr Met Asn Gly  
 1250 1255 1260  
 Thr Asn Ile Leu His Arg Gly Ala Tyr Tyr Val Leu Lys Asp Trp Ala  
 265 1270 1275 1280  
 Thr Asn Gln Tyr Phe Asn Ile Ala Lys Thr Asn Glu Val Phe Leu Pro  
 1285 1290 1295  
 Leu Gln Leu Gln Asn Lys Asp Ala Gln Thr Gly Phe Ile Ser Asp Ala  
 1300 1305 1310

16/36

Ser Gly Val Lys Tyr Tyr Ser Ile Ser Gly Tyr Gln Ala Lys Asp Thr  
 1315 1320 1325  
 Phe Ile Glu Asp Gly Asn Gly Asn Trp Tyr Tyr Phe Asp Lys Asp Gly  
 1330 1335 1340  
 Tyr Met Val Arg Ser Gln Gln Gly Glu Asn Pro Ile Arg Thr Val Glu  
 345 1350 1355 1360  
 Thr Ser Val Asn Thr Arg Asn Gly Asn Tyr Tyr Phe Met Pro Asn Gly  
 1365 1370 1375  
 Val Glu Leu Arg Lys Gly Phe Gly Thr Asp Asn Ser Gly Asn Val Tyr  
 1380 1385 1390  
 Tyr Phe Asp Asp Gln Gly Lys Met Val Arg Asp Lys Tyr Ile Asn Asp  
 1395 1400 1405  
 Asp Ala Asn Asn Phe Tyr His Leu Asn Val Asp Gly Thr Met Ser Arg  
 1410 1415 1420  
 Gly Leu Phe Lys Phe Asp Ser Asp Thr Leu Gln Tyr Phe Ala Ser Asn  
 425 1430 1435 1440  
 Gly Val Gln Ile Lys Asp Ser Tyr Ala Lys Asp Ser Lys Gly Asn Lys  
 1445 1450 1455  
 Tyr Tyr Phe Asp Ser Ala Thr Gly Asn Asn Asp Thr Gly Lys Ala Gln  
 1460 1465 1470  
 Thr Trp Asp Gly Asn Gly Tyr Tyr Ile Thr Ile Asp Ser Asp Ala Asn  
 1475 1480 1485  
 Asn Thr Ile Gly Val Asn Thr Asp Tyr Thr Ala Tyr Ile Thr Ser Ser  
 1490 1495 1500  
 Leu Arg Glu Asp Gly Leu Phe Ala Asn Ala Pro Tyr Gly Val Val Thr  
 505 1510 1515 1520  
 Lys Asp Gln Asn Gly Asn Asp Leu Lys Trp Gln Tyr Ile Asn His Thr  
 1525 1530 1535  
 Lys Gln Tyr Glu Gly Gln Gln Val Gln Val Thr Arg Gln Tyr Thr Asp  
 1540 1545 1550  
 Ser Lys Gly Val Ser Trp Asn Leu Ile Thr Phe Ala Gly Gly Asp Leu  
 1555 1560 1565  
 Gln Gly Gln Arg Leu Trp Val Asp Ser Arg Ala Leu Thr Met Thr Pro  
 1570 1575 1580  
 Phe Lys Thr Met Asn Gln Ile Ser Phe Ile Ser Tyr Ala Asn Arg Asn  
 585 1590 1595 1600  
 Asp Gly Leu Phe Leu Asn Ala Pro Tyr Gln Val Lys Gly Tyr Gln Leu  
 1605 1610 1615  
 Ala Gly Met Ser Asn Gln Tyr Lys Gly Gln Gln Val Thr Ile Ala Gly  
 1620 1625 1630

17/36

Val Ala Asn Val Ser Gly Lys Asp Trp Ser Leu Ile Ser Phe Asn Gly  
 1635 1640 1645  
 Thr Gln Tyr Trp Ile Asp Ser Gln Ala Leu Asn Thr Asn Phe Thr His  
 1650 1655 1660  
 Asp Met Asn Gln Lys Val Phe Val Asn Thr Thr Ser Asn Leu Asp Gly  
 665 1670 1675 1680  
 Leu Phe Leu Asn Ala Pro Tyr Arg Gln Pro Gly Tyr Lys Leu Ala Gly  
 1685 1690 1695  
 Leu Ala Lys Asn Tyr Asn Asn Gln Thr Val Thr Val Ser Gln Gln Tyr  
 1700 1705 1710  
 Phe Asp Asp Gln Gly Thr Val Trp Ser Gln Val Val Leu Gly Gly Gln  
 1715 1720 1725  
 Thr Val Trp Val Asp Asn His Ala Leu Ala Gln Met Gln Val Ser Asp  
 1730 1735 1740  
 Thr Asp Gln Gln Leu Tyr Val Asn Ser Asn Gly Arg Asn Asp Gly Leu  
 745 1750 1755 1760  
 Phe Leu Asn Ala Pro Tyr Arg Gly Gln Gly Ser Gln Leu Ile Gly Met  
 1765 1770 1775  
 Thr Ala Asp Tyr Asn Gly Gln His Val Gln Val Thr Lys Gln Gly Gln  
 1780 1785 1790  
 Asp Ala Tyr Gly Ala Gln Trp Arg Leu Ile Thr Leu Asn Asn Gln Gln  
 1795 1800 1805  
 Val Trp Val Asp Ser Arg Ala Leu Ser Thr Thr Ile Met Gln Ala Met  
 1810 1815 1820  
 Asn Asp Asn Met Tyr Val Asn Ser Ser Gln Arg Thr Asp Gly Leu Trp  
 825 1830 1835 1840  
 Leu Asn Ala Pro Tyr Thr Met Ser Gly Ala Lys Trp Ala Gly Asp Thr  
 1845 1850 1855  
 Arg Ser Ala Asn Gly Arg Tyr Val His Ile Ser Lys Ala Tyr Ser Asn  
 1860 1865 1870  
 Glu Val Gly Asn Thr Tyr Tyr Leu Thr Asn Leu Asn Gly Gln Ser Thr  
 1875 1880 1885  
 Trp Ile Asp Lys Arg Ala Phe Thr Val Thr Phe Asp Gln Val Val Ala  
 1890 1895 1900  
 Leu Asn Ala Thr Ile Val Ala Arg Gln Arg Pro Asp Gly Met Phe Lys  
 905 1910 1915 1920  
 Thr Ala Pro Tyr Gly Glu Ala Gly Ala Gln Phe Val Asp Tyr Val Thr  
 1925 1930 1935  
 Asn Tyr Asn Gln Gln Thr Val Pro Val Thr Lys Gln His Ser Asp Ala  
 1940 1945 1950

18/36

Gln Gly Asn Gln Trp Tyr Leu Ala Thr Val Asn Gly Thr Gln Tyr Trp  
 1955 1960 1965

Ile Asp Gln Arg Ser Phe Ser Pro Val Val Thr Lys Val Val Asp Tyr  
 1970 1975 1980

Gln Ala Lys Ile Val Pro Arg Thr Thr Arg Asp Gly Val Phe Ser Gly  
 985 1990 1995 2000

Ala Pro Tyr Gly Glu Val Asn Ala Lys Leu Val Asn Met Ala Thr Ala  
 2005 2010 2015

Tyr Gln Asn Gln Val Val His Ala Thr Gly Glu Tyr Thr Asn Ala Ser  
 2020 2025 2030

Gly Ile Thr Trp Ser Gln Phe Ala Leu Ser Gly Gln Glu Asp Lys Leu  
 2035 2040 2045

Trp Ile Asp Lys Arg Ala Leu Gln Ala  
 2050 2055

<210> 3  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: artificial  
 sequence

<400> 3  
 actgcggccg catgccattt acagaaaaag 30

<210> 4  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: artificial  
 sequence

<400> 4  
 actgctcgag ttatgctgac acagcatttc 30

<210> 5  
 <211> 7  
 <212> PRT  
 <213> Leuconostoc mesenteroides

<400> 5  
 Lys Thr Asn Glu Val Phe Leu  
 1 5

<210> 6

19/36

<211> 7  
<212> PRT  
<213> Leuconostoc mesenteroides

<400> 6  
Lys Phe Asp Ser Asp Arg Leu  
1 5

<210> 7  
<211> 19  
<212> PRT  
<213> Leuconostoc mesenteroides

<400> 7  
Lys Asn Met Gln Asp Thr Phe Thr Phe Asp Gln Leu Ala Gln Gly Met  
1 5 10 15

Glu Phe Tyr

<210> 8  
<211> 15  
<212> PRT  
<213> Leuconostoc mesenteroides

<400> 8  
Lys Ser Thr Glu Trp Leu Arg Asp Ala Ile Asp Leu Phe Val Lys  
1 5 10 15

<210> 9  
<211> 22  
<212> PRT  
<213> Leuconostoc mesenteroides

<400> 9  
Lys Gly Ser Glu Phe Leu Leu Ala Asn Asp Ile Asp Asn Ser Asn Pro  
1 5 10 15

Ile Val Gln Ala Glu Gln  
20

<210> 10  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 10  
ytgrtcraan gtraangtrt cytgcatrtt ytt

33

<210> 11  
<211> 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial sequence

&lt;400&gt; 11

gaygcnathg ayytnttygt naar

24

&lt;210&gt; 12

&lt;211&gt; 837

&lt;212&gt; DNA

&lt;213&gt; Leuconostoc mesenteroides

&lt;400&gt; 12

```

gatgcaattg atagttttgt gaagacgcaa gctaattgga ataagcaaac agaagatgaa 60
gctttcgatg gtttgcagtg gcttcaaggg ggattcctag cttatcaaga tgattcacat 120
cggacgccga atactgattc aggaaataac agaaaactag gacgtcaacc aattaatatc 180
gatggttcga aagatacaac tgatggtaaa ggctctgaat tcttattagc taacgatatt 240
gacaactcaa atccgattgt tcaagctgag caattaaact ggctacacta tttaatgaat 300
tttggtagta ttacaggtaa taatgacaat gcgaattttg atggcattcg tgtagatgct 360
gttgataatg ttgatgctga tttactaaaa atagctggcg attattttta agctctatat 420
ggtagacata aaagcgacgc caatgccaat aagcattttg ctatttttaga agactggaac 480
ggtaaagatc ctcagtatgt taatcaacag ggcaatgcmc aattaacaat ggattacaca 540
gttacttcac agtttggcaa ttctctaaca catggcgcca acaacaggag taacatgtgg 600
tatttcttag atactggcta ttatcttaat ggagatctta ataagaagat agtagataag 660
aaccgtccaa attctggcac tttggttaac agaattgcta attcaggtga taaaaaagtt 720
attccaaatt atagttttgt tagagcacat gattacgatg ctcaagatcc aattagaaaa 780
gccatgattg atcatggtat tattaataaac atgcaggata ctttcacttt tgaccaa 837

```

&lt;210&gt; 13

&lt;211&gt; 2862

&lt;212&gt; DNA

&lt;213&gt; Leuconostoc mesenteroides

&lt;400&gt; 13

```

gatcaaatag atgctaccgt tatacgagac gttaaggatt atcccaatct agtcgttttg 60
cgtaccttgt cgaaagctta tggctcggca aacctgagaa tcggctatgg cgtcatgcag 120
gaaccacttt atcaggttat gcaggccgta cgtttaccat ataattttaa tacctatcaa 180
atcacagggtg cagtagctgc ccttagtgat caactttatc tgcaatcagt tgttgctaag 240
gtgaagtctg aacgtgaaaa atttgaacaa tttttgacga aacaccagtt taagtattat 300
caatcacaaa ccaactttct ctggattaaa gttgggtgatg cgaaacgtgt tggtagggct 360
cttctgtcag aagggtatca aattaatgac cgcctaaatg ccgaatggat tcgcattgca 420
ttaggaactg tgtctgataa tgaggggatg cagcgcattt tattgaattg ttaaaaaata 480
gctaagagag tatgttcttc tcttacctat ttttatttgt aattcctatt atttaatttt 540
gcatgacaat attaatagcg tgttacgatt ctactattta atgtaataaa aattaataaa 600
tatggtatta tcttatatgg gtgatagatg caccaaatac tgtatcatgt ctggtcacat 660
gaaagggaga ataattaatg aaacaacaag aaacagttac ccgtaaaaaa ctttataaat 720
ccggtaagggt ttgggttgca gcagctactg catttgcggt attgggggtt tcaactgtaa 780
caacagtcca tgcggataca aattcgaatg tcgctgttaa gcaaataaat aatacaggaa 840
ccaatgattc tggcgaaaaa aaggatccgg ttccatcaac taataatgat agtttgaagc 900
aaggaacaga tggtttttgg tatgattcag acggcaatcg tgtcgatcag aagaccaatc 960
agattctgct tactgcggaa caacttaaaa aaaataacga aaaaaattta tcagtaatca 1020
gtgatgatac atcaaaaaaa gatgatgaaa atatttctaa gcagaccaa attgctaata 1080
aacaacagat agatactgct aaaggcctga ctaccagtaa tttatctgat cccatcactg 1140
ggggtcacta tgaaaatcac aatggctact ttgtttatat agatgcttca ggaaaacaag 1200
taacagggtt gcaaaaatatt gatggtaatt tacaatattt tgatgacaat ggatatcaag 1260

```



tcaagggatc	cttccgagat	gtcaacggca	agcatatcta	ttttgattca	gtaacagggga	1320
aagctagttc	aaatgttgat	attgttaacg	gtaaagctca	aggatatgat	gcgcaaggca	1380
accaattaaa	gaaaagttat	gtcgccgata	gttctgggca	aacttactat	tttgatggta	1440
atggccaacc	gttaatcggc	ttgcaaaca	ttgatgggaa	cctacaatat	tttaaccaac	1500
aaggggttca	aataaagggt	ggtttccaag	atgttaacaa	taaacgtatt	tattttgcac	1560
caaacacagg	taatgccgtt	gccaatactg	aaataattaa	cggtaaatta	caggggctg	1620
acgcaaattg	taaccaggta	aagaatgcat	ttagtaaaaga	tggtgcagga	aatacatttt	1680
attttgacgc	aaacggtgtg	atgttaacag	ggttgcaaac	tatttcagga	aagacatatt	1740
atcttgatga	acaaggacac	ctgagaaaaa	attacgcggg	aacattcaat	aatcagttta	1800
tgtacttcga	tgctgatata	ggtgcgggta	aaacagcgat	tgaatatcaa	tttgatcaag	1860
gattggatc	acaaagtaat	gaaaatactc	ctcacaatgc	cgcaaagtct	tatgataaaa	1920
gtagttttga	aaatgttgat	ggttacttaa	cagcagatac	atggtatcgt	ccaaccgata	1980
ttttaaaaaa	tgagataact	tggaacggcat	ctaccgaaac	tgatatgcgt	ccgcttttaa	2040
tgacatgggtg	gcctgacaaa	caaacacaag	caaattactt	gaattttatg	tctagtaaag	2100
gacttggtat	aacgaccact	tatacagcag	ctacgtcaca	aaaaacacta	aatgacgcag	2160
cctttgttat	tcaaacagca	attgaacaac	aaatatcttt	gaaaaaaagt	actgagtgg	2220
tacgtgatgc	aattgatagt	tttgtgaaga	cgcaagctaa	ttggaataag	caaacagaag	2280
atgaagcttt	cgatggtttg	cagtggcttc	aagggggatt	cctagcttat	caagatgatt	2340
cacatcggac	gccgaatact	gattcaggaa	ataacagaaa	actaggacgt	caaccaatta	2400
atatcgatgg	ttcgaaagat	acaactgatg	gtaaaggctc	tgaattctta	ttagctaacg	2460
atattgacaa	ctcaaatccg	attgttcaag	ctgagcaatt	aaactggcta	cactatttaa	2520
tgaatttttg	tagtattaca	ggtaataatg	acaatgcgaa	ttttgatggc	attcgtgtag	2580
atgctgttga	taatgttgat	gctgatttac	taaaaatagc	tggcgattat	tttaaagctc	2640
tatatggtac	agataaaagc	gacgccaatg	ccaataagca	tttgtctatt	ttagaagact	2700
ggaacggtaa	agatcctcag	tatgttaatc	aacagggcaa	tgcgcaatta	acaatggatt	2760
acacagttac	ttcacagttt	ggcaattctc	taacacatgg	cgccaacaac	aggagtaaca	2820
tgtggtat	cttagatact	ggctattatc	ttaatggaga	tc		2862

&lt;210&gt; 14

&lt;211&gt; 3970

&lt;212&gt; DNA

<213> *Leuconostoc mesenteroides*

&lt;400&gt; 14

gatcaaggat	tggtatcaca	aagtaatgaa	aatactctc	acaatgccgc	aaagtcttat	60
gataaaagta	gttttgaaaa	tggtgatgg	tacttaacag	cagatacatg	gtatcgctcca	120
accgatattt	taaaaaatgg	agatacttgg	acggcatcta	ccgaaactga	tatgctgccg	180
cttttaatga	catggtggcc	tgacaaaca	acacaagcaa	attacttgaa	ttttatgtct	240
agtaaaggac	ttggtataac	gaccacttat	acagcagcta	cgtcacaaaa	aacactaaat	300
gacgcagcct	ttgttattca	aacagcaatt	gaacaacaaa	tatctttgaa	aaaaagtact	360
gagtggttac	gtgatgcaat	tgatagtttt	gtgaagacgc	aagctaattg	gaataagcaa	420
acagaagatg	aagctttcga	tggtttgcag	tggcttcaag	ggggattcct	agcttatcaa	480
gatgattcac	atcggacgcc	gaatactgat	tcaggaaata	acagaaaact	aggacgtcaa	540
ccaattaata	tcgatgggtc	gaaagataca	actgatggta	aaggctctga	attcttatta	600
gctaacgata	ttgacaactc	aaatccgatt	gttcaagctg	agcaattaaa	ctggctacac	660
tattttaatga	attttggtag	tattacaggt	aataatgaca	atgcgaattt	tgatggcatt	720
cgtgtagatg	ctggtgataa	tggtgatgct	gatttactaa	aaatagctgg	cgattatttt	780
aaagctctat	atggtacaga	taaaagcgac	gccaatgcc	ataagcattt	gtctatttta	840
gaagactgga	acggtaaaga	tcctcagtat	gttaatcaac	agggcaatgc	gcaattaaca	900
atggattaca	cagttacttc	acagtttggc	aattctctaa	cacatggcgc	caacaacagg	960
agtaacatgt	ggtatttctt	agatactggc	tattatctta	atggagatct	taataagaag	1020
atagtagata	agaaccgtcc	aaattctggc	actttgggta	acagaattgc	taattcaggt	1080
gatacaaaaag	ttattccaaa	ttatagtttt	gtagagcac	atgattacga	tgctcaagat	1140
ccaattagaa	aagccatgat	tgatcatggt	attattaaaa	acatgcagga	tactttcact	1200
tttgaccaac	tggtcaggg	aatggaattc	tactataaag	atcaagagaa	tccgtctgg	1260
ttcaaaaagt	ataacgatta	taacttacct	agtcttatg	caatgttggt	gactaataag	1320
gatactgtac	ctcgtgtcta	ttatggagat	atgtacctcg	aaggcgggca	atatatggaa	1380
aaagggacga	tttacaatcc	tgctatttca	gcgttgctca	aagctagaat	aaaatatggt	1440
tctggtgggc	aaacaatggc	taccgatagt	tctggaaaag	accttaaaga	tggcgaaact	1500

gatttggttaa	caagtgttcg	atttggtaaa	ggaattatga	catcagatca	aaccacaaca	1560
caagacaata	gccaaagatta	taaaaatcaa	ggcatcggtg	tcattgttgg	taataaccct	1620
gaccttaagt	tgaacaatga	taagaccatt	accttgcata	tgggaaaggc	gcataagaat	1680
caactttacc	gtgccttagt	attatcaa	gactcaggaa	ttgatgttta	tgatagtgat	1740
gataaagcac	caactttgag	aacaaatgac	aacggtgact	tgattttcca	taagacaaat	1800
acgtttgtga	agcaagatgg	aactattata	aattacgaaa	tgaagggatc	attaaatgct	1860
ttaatttcag	gttatttagg	tgtctgggtg	ccagttggag	ctagtgattc	acaagatgct	1920
cgtacagtgg	caactgagtc	atcatcaagt	aatgatgggt	ctgtattcca	ttcaaatgct	1980
gcattagatt	ctaagtgtat	atatgaaggc	ttttcaaact	ttcaagcgat	gccgacttct	2040
cctgagcaaa	gtacaaaatg	tgttattgca	acaaaggcta	acttatttaa	agaattaggt	2100
attactagtt	ttgagttagc	acctcaatat	aggtctagt	gtgacactaa	ttacgggtggc	2160
atgtcattct	tagattcttt	cttaataaat	ggttatgcat	ttaccgatag	atatgattta	2220
ggctttaaca	aagcagacgg	gaatcctaac	ccaacaaagt	atggaacaga	tcaagattta	2280
cgtaatgcaa	tagaggcatt	acacaaaaac	ggcatgcagg	ctatagctga	ttgggttcc	2340
gaccaaata	atgctttacc	aggaaaggaa	gttgttaccg	ctactagagt	agacgaacgg	2400
ggaaatcaac	taaaagacac	agattttgtc	aacttactct	atgttgctaa	tactaaaagt	2460
agtgggtgtg	attatcaggc	aaagtatggc	ggcgaatttt	tagataaatt	aagagaagag	2520
tacccatcgt	tattcaaaca	gaaccaagta	tcgacaggtc	agccaattga	tgcttctaca	2580
aaaattaagc	aatggtcagc	taaatatatg	aatgggacca	atattttaca	tcgaggtgct	2640
tattatgttt	tgaagactg	ggctactaac	cagtatttta	acattgcaaa	aacgaatgaa	2700
gtatttttgc	cactacagtt	gcagaataaa	gatgcgcaaa	ctggtttcat	tagtgatgcc	2760
tccggtgtaa	aatattactc	aattagtggt	tatcaagcaa	aagatacttt	tattgaagat	2820
ggtaatggga	attgggtatta	ctttgataaa	gatgggtaca	tgggtgcgtc	gcagcaagga	2880
gaaaatccta	taagaacagt	cgaaactagt	gtcaacacac	gaaacggtaa	ttattacttt	2940
atgccaaatg	gtgtcgagtt	gcgcaaaggc	tttggaaagg	ataatagtg	taatgtctat	3000
tattttgatg	atcaaggtaa	gatgggtgag	gataaatata	ttaacgatga	tgctaataat	3060
ttttatcact	taaatgttga	tgggactatg	tctcgaggac	tatttaaatt	tgattctgat	3120
actctacagt	attttgctag	taatggtgtc	caaataaaag	atagttatgc	gaaggatagt	3180
aaaggcaata	aatattat	tgactcagct	acaggaaata	acgatactgg	gaaagcccaa	3240
acttgggatg	gtaatggcta	ctatatctat	attgattctg	atgcgaacaa	tacaattggg	3300
gtaaacacag	actacactgc	ctacatcact	agctcgctgc	gcgaagatgg	cttatttgct	3360
aacgcacctt	acggtgttgt	aacaaaagac	caaaatggta	acgatcttaa	gtggcagtat	3420
attaaccata	cgaacagta	cgaagggcaa	caagtgcag	tcacgcgtca	atacacagac	3480
agtaaggag	tcagctggaa	cttaattacc	tttgctgggtg	gtgatttaca	aggacaaagg	3540
ctttgggtgg	atagtcgtgc	gttaactatg	acaccattta	aaacgatgaa	ccaaataagc	3600
ttcattagtt	atgctaaccg	caatgatggg	ttgtttttga	atgcgccata	ccaagtcaag	3660
gggtatcaat	tagctgggat	gtccaaccaa	tacaagggcc	aacaagtgc	cattgctggg	3720
gtggcgaaag	tttctggaaa	agactggagt	ctgattagtt	ttaatgggac	acagtactgg	3780
attgatagtc	aggcattgaa	taccaatttc	acacatgaca	tgaacaaaa	ggtctttgtc	3840
aatacaacta	gtaatcttga	tgggttat	ttaaatgcgc	cataccgtca	accgggttat	3900
aagttagccg	gtttggctaa	aaattacaac	aaccaaagg	ttactgttag	tcaacagtac	3960
tttgatgatc						3970

&lt;210&gt; 15

&lt;211&gt; 1917

&lt;212&gt; DNA

<213> *Leuconostoc mesenteroides*

&lt;400&gt; 15

gatcaaggta	agatgggtgag	agataaatac	attaacgatg	atgctaataa	tttttatcac	60
ttaaatgttg	atgggactat	gtctcgagga	ctatttaaat	ttgattctga	tactctacag	120
tattttgtcta	gtaattggtg	ccaaataaaa	gatagttatg	cgaaggatag	taaaggcaat	180
aaatattatt	ttgactcagc	tacaggaaat	aacgatactg	ggaaagccca	aacttgggat	240
ggtaattggct	actataattac	tattgattct	gatgcgaaca	atacaattgg	ggttaacaca	300
gactacactg	cctacatcac	tagctcgctg	cgcgaaagatg	gcttatttgc	taacgcacct	360
tacgggtgttg	taacaaaaga	ccaaaatggt	aacgatctta	agtggcagta	tattaaccat	420
acgaaacagt	acgaagggca	acaagtgcaa	gtcacgcgtc	aatacacaga	cagtaaggga	480
gtcagctgga	acttaattac	ctttgctgggt	ggtgatttac	aaggacaaag	gctttgggtg	540
gatagtcgtg	cgtaactat	gacaccattt	aaaacgatga	accaaataag	cttcattagt	600

```

tatgctaacc gcaatgatgg gttgtttttg aatgcgccat accaagtcaa ggggtatcaa 660
ttagctggga tgtccaacca atacaagggc caacaagtga ccattgctgg ggtggcgaac 720
gtttctggaa aagactggag tctgattagt tttaatggga cacagtactg gattgatagt 780
caggcattga ataccaattt cacacatgac atgaaccaa aggtctttgt caatacaact 840
agtaatcttg atgggttatt cttaaatgcg ccataccgtc aaccgggtta taagttagcc 900
ggtttggtta aaaattacaa caaccaaacg gttactgtta gtcaacagta ctttgatgat 960
caaggcacgg tctggagtca ggttgtcctt gggggtcaga cggctctggg tgataaccat 1020
gcattggcac agatgcaagt tagtgatata gaccaacagc tctatgtgaa tagcaatggg 1080
cggaatgatg ggttatttctt gaatgcgcca tatcgtgggc aagggtcaca actgataggc 1140
atgacggcag attataatgg gcaacatgta caagtgacca agcaagggca agatgcctat 1200
ggtgcacaat ggcgtcttat tacgctaaat aatcaacagg tctgggttga tagtcgcgct 1260
ttgagcacia caatcatgca agccatgaat gataatatgt atgtaaatag cagccaacgg 1320
acagatggct tgtgggttaa cgcaccttat acgatgagtg gggctaaatg ggctgggtgat 1380
acacgttcag ctaatgggag ctatgtccat atttcaaaag cttattcaaa cgaagtccgg 1440
aatacatatt acttgacgaa tttgaatggt caaagcacat ggattgacaa gcgggctgtt 1500
actgtgacct tcgatcagg ggtggcatta aatgcaacga ttgtggcacg ccaacgacca 1560
gatgggatgt ttaagacagc accatatggt gaagcggggg cgcagtttgt cgattatgtg 1620
acaaactata accagcaaac cgtgccagta acaaagcaac attcagatgc tcaggggaat 1680
caatggtact tagcgacagt gaatgggaca caatactgga ttgatcaacg gtcattttca 1740
ccagtagtaa cgaaggtggt tgattatcaa gctaagattg tgccacggac aacacgtgat 1800
ggtgtgttta gtggcgccac ctatggggaa gtgaatgcta agctagttaa catggcaact 1860
gcgtatcaaa atcaagttgt ccatgcgaca ggggaatata cgaatgcttc agggatc 1917

```

&lt;210&gt; 16

&lt;211&gt; 4066

&lt;212&gt; DNA

<213> *Leuconostoc mesenteroides*

&lt;400&gt; 16

```

gatcttaagt ggcagtatat taaccatacg aaacagtagc aagggaaca agtgcaagtc 60
acgcgtcaat acacagacag taagggagtc agctggaact taattacctt tgctgggtgt 120
gatttacaag gacaaaggct ttgggtggat agtcgtgcgt taactatgac accatttaaa 180
acgatgaacc aaataagctt cattagttat gctaaccgca atgatgggtt gtttttgaat 240
gcgccatacc aagtcaagggt gtatcaatta gctgggatgt ccaaccaata caagggcaa 300
caagtgacca ttgctggggt ggogaacgtt tctggaaaag actggagtct gattagtttt 360
aatgggacac agtactggtt tgatagtcag gcattgaata ccaatttcac acatgacatg 420
aaccaaaagg tctttgtcaa tacaactagt aatcttgatg ggttattctt aaatgcgcca 480
taccgtcaac cgggttataa gttagccggt ttggctaaaa attacaacaa ccaaacggtt 540
actgttagtc aacagttact tgatgatcaa ggcacggctt ggagtcaggt tgcctttggg 600
ggtcagacgg tctgggttga taaccatgca ttggcacaga tgcaagttag tgatacagac 660
caacagctct atgtgaatag caatggctcg aatgatgggt tattcttgaa tgcgccatat 720
cgtgggtcaag ggtcacaact gataggcatg acggcagatt ataatgggca acatgtacaa 780
gtgaccaagc aagggaaga tgacctatgg gcacaatggc gtcttattac gctaaataat 840
caacaggtct ggggttgatag tcgcgctttg agcacaacaa tcatgcaagc catgaatgat 900
aatatgtatg taaatagcag ccaacggaca gatggcttgt ggttaaacgc accttatacg 960
atgagtgggg cttaaagggt tggatgata cgttcagcta atgggcgcta tgtccatatt 1020
tcaaaagctt attcaaacga agtcggcaat acatattact tgacgaattt gaatggtcaa 1080
agcacatgga ttgacaagcg ggcgtttact gtgaccttcg atcaggtggt ggcattaaat 1140
gcaacgattg tggcacgcca acgaccagat gggatgttta agacagcacc atatggtgaa 1200
gcggggggcg agtttgtcga ttatgtgaca aactataacc agcaaaccgt gccagtaaca 1260
aagcaacatt cagatgctca ggggaatcaa tgggtacttag cgacagtga tgggacacaa 1320
tactggattg atcaacggtc attttcacca gtagtaacga aggtggttga ttatcaagct 1380
aagatttgtc cacggacaac acgtgatggt gtgttttagtg gcgcacccta tggggaagtg 1440
aatgctaagc tagttaacat ggcaactgcg tatcaaaatc aagttgtcca tgcgacaggg 1500
gaatatacga atgcttcagg gatcacatgg agtcagttcg cgtaagcgg gcaagaagac 1560
aagctatgga ttgataagcg tgctttgcaa gcttaaggga aggattcgac aaaggagggt 1620
aacattatca gcggatggtg ttatcctcct ttctgtact cagtatttcc caaataattg 1680
agacagtttc atgacaaatc aacaaaacta gtgtcaatgc ctcggttatg gggtaaacta 1740
ctattagtta aagggttgtt gcataataat atcacattca atatattatg tatttttatc 1800

```

```

tgattatgtg attttttgag atttggagcg aaaatgaata aagtaataat taataaaaagt 1860
ttctgtgttt tggtaatgag tttgttgctg atatttctat tctcattaag agtagacgct 1920
cgttccaata gtggctacaa tcaaatactt tttaatcaac atgaaattgc ttatgcgct 1980
tttagtcaag taccttggtg tgcaactgtg aatgtcggaa tgtctgcaga caaaaaataat 2040
atctatacag ctattgatat ggggccatag tatcaaggaa cttctttttt tccgtggtgg 2100
tatcaacatg ataattatga ttatcatgat ggttatcaac ttgcgattga caatcaatcg 2160
cttttcatag aaatggttga tacaggaagg tcgtcgtttg atgttattga gttgtataca 2220
ggtaaaaaag ttggttatgg aaaattcaca gcctataact ataacaagac ttcatggcag 2280
tttcaaatat ctttaaaggc gataaagaaa attttaaata ataacatatc agaaaatgcg 2340
acaattagtt tgtacaatgg ttatctattc aaacaagccg aaagtatcac gtatgctggt 2400
tcaccaacag ggccatggtt acttgccggtt attggtttgg tttttgctgg tgggggctat 2460
tacctgcaaa atcagcgccg aaaaacgatt acggcatcat caacaaacga ggtaacccat 2520
gcttaataat gctgttatta ttggtctaatt tattatttgg gtctatgtat tatctgttct 2580
gaagcgtgct aaaacggatg cgttttattt ttggctggc agcgcaggct tgttcgtcat 2640
tcttctttta tggagtaaac cttatggtgt ttggttgtt tcgaccattc tcacttggtc 2700
ggccgggggtt gttggtcatc tcactggctt atttgatact ttctatgctg cactgttat 2760
tcaggttgtg gcgaatcatc ataccagcat tttattgggt gattatgaat gttcaggcat 2820
tattgaacaa acggcatttt ggggattaat tgccttttat cctgtgtatg atgcacaaaa 2880
gcggttaatg ttggcgctta ttggggcatt atgggttttc cttgcaaata cgttacgact 2940
atcttttgtt gcagttgtaa tttattattt tggcgatggc gctttttatg ttgctcattc 3000
aattatcgga cgcctgttgt tctattcaat tgtaatcttg ctttattatc tagtcttcat 3060
aaaggggcaa cttgttaatc aaatgctcaa aaaacaggga taaactatgt cattttattt 3120
acatcaattt acttatcagt taggtttttg gttcacttgg ttgctcattc cgtttgttgt 3180
ggaaattttc ccagctatta tttcaatcta gttttgataa aaacatcaaa aaagcatcaa 3240
gtgatgcaag aaccattgaa attaccaatg gtatctattg tattacctat atataattca 3300
ggacaaacat tataccagtg tattcagtcg attagtcaat caacctatcc aaaacaatta 3360
atccaaatta ttgccgtgaa taatcaaagc actgataaca gttttacagt gtttaatcag 3420
gctcaagctg atttcccaat acttagaatg cagtggatga acacagatca aggtaaagcg 3480
cgtgcactaa atgctgctat ttataatagt atggggcaat atattattaa tttggatact 3540
gatggttggg tagaacccaa tgccttaaaa cggtttgtgc tttattttga aaatcactca 3600
gaaattgatg tagcaactgg tacgattctg acacaaaaaa aaatgattca aaaaacgcaa 3660
agtaaattggc ttaaattgct acagttaaac gaatattttg agtatgcaca atcattctta 3720
tcaggccgta gtattgaaaa ccgtggtaat cgtttgttca caatgtcagg tgcattttca 3780
gcgttttagac gtgatgtatt agttcagaca tttatgtata atgttgacac ggttggtgaa 3840
gatactgata tgacatttca actgcgattc cgcctaggca agcgtattgg tttctgtgat 3900
gatgccatgt tttatgttga accaatatca ggttatagtg aactttattt acaacggcag 3960
cgttggcagc gtgggcaaat tgaagtcgca caaaatttca tgcagaataa gttgagtgtc 4020
cgtcagattt ttactaactt tatgattagt cgattaatga ttgatc 4066

```

&lt;210&gt; 17

&lt;211&gt; 7387

&lt;212&gt; DNA

<213> *Leuconostoc mesenteroides*

&lt;400&gt; 17

```

gatcaaatag atgctaccgt tatacgagac gttaaggatt atcccaatct agtcgttttg 60
cgtaccttgt cgaaagctta tggctctggca aacctgagaa tcggctatgg cgtcatgcag 120
gaaccacttt atcaggttat gcaggccgta cgtttaccat ataattttaa tacctatcaa 180
atcacagggtg cagttagctgc ctttagtgat caactttatc tgcaatcagt tgttgctaag 240
gtgaagtctg aacgtgaaaa atttgaacaa tttttgacga aacaccagtt taagtattat 300
caatcacaaa ccaactttct ctggattaaa gttggtgatg cgaaacgtgt tgggtagggt 360
cttctgtcag aagggatatca aattaatgac cgcctaaatg ccgaatggat tcgcattgca 420
ttaggaactg tgtctgataa tgaggggatg cagcgcattt tattgaattg ttaaaaaata 480
gctaagagag tatgttcttc tcttacctat ttttatttgt aattcctatt atttaatttt 540
gcatgacaat attaatagcg tgttacgatt ctactattta atgttaataa aattaataaa 600
tatggtatta tcttatatgg gtgatagatg caccaaatac tgtatcatgt ctggtcacat 660
gaaagggaga ataattaatg aaacaacaag aaacagttac ccgtaaaaaa ctttataaat 720
ccggttaaggt ttgggttgca gcagctactg catttgcggt attgggggtt tcaactgtaa 780
caacagttcca tgcggatata aattcgaatg tcgctgttaa gcaaataaat aatacaggaa 840

```

```

ccaatgattc tggcgaaaaa aaggtaccgg ttccatcaac taataatgat agtttgaagc 900
aaggaacaga tgggttttgg tatgattcag acggcaatcg tgtcgatcag aagaccaatc 960
agattctgct tactgcgga caacttaaaa aaaataacga aaaaaattta tcagtaatca 1020
gtgatgatac atcaaaaaaa gatgatgaaa atatttctaa gcagaccaa attgctaate 1080
aacaacagat agatactgct aaaggcctga ctaccagtaa tttatctgat cccatcactg 1140
ggggtcacta tgaaaatcac aatggctact ttgtttatat agatgcttca ggaaaacaag 1200
taacagggtt gcaaaatatt gatggtaatt tacaatat ttgtgattca gtaacaggga 1320
tcaagggatc cttccgagat gtcaacggca agcatatcta ttttgattca gtaacaggga 1320
aagctagttc aaatgttgat attgttaacg gtaaagctca aggatattgat gcgcaaggca 1380
accaattaaa gaaaagttat gtcgccgata gttctgggca aacttactat tttgatggta 1440
atggccaacc gttaatcggc ttgcaaaaca ttgatgggaa cctacaatat tttaaccaac 1500
aaggggttca aataaagggt gggttccaag atgttaacaa taaacgtatt tattttgcac 1560
caaacacagg taatgccgtt gccaatactg aaataattaa cggtaaatta caggggcgtg 1620
acgcaaatgg taaccaggta aagaatgcat ttagtaaaga tgttgcagga aatacatatt 1680
attttgacgc aaacgggtgt atgttaacag gggtgcaaac tatttcagga aagacatatt 1740
atcttgatga acaaggacac ctgagaaaaa attacgcggg aacattcaat aatcagttta 1800
tgtacttcca tgctgataca ggtgcgggta aaacagcgat tgaatatcaa tttgatcaag 1860
gattgggtatc acaaagtaat gaaaatactc ctcaaatgc cgcgaagtct tatgataaaa 1920
gtagttttga aaatgttgat gggtacttaa cagcagatac atggtatcgt ccaaccgata 1980
ttttaaaaaa tggagatact tggacggcat ctaccgaaac tgatatgcgt ccgcttttaa 2040
tgacatgggtg gcctgacaaa caaacacaag caaattactt gaattttatg tctagtaaag 2100
gacttggtat aacgaccact tatacagcag ctacgtcaca aaaaacacta aatgacgcag 2160
cctttgttat tcaaacagca attgaacaac aaatatcttt gaaaaaaagt actgagtgg 2220
tacgtgatgc aattgatagt tttgtgaaga cgcaagctaa ttggaataag caaacagaag 2280
atgaagcttt cgatggtttg cagtggcttc aagggggatt cctagcttat caagatgatt 2340
cacatcggac gccgaatact gattcaggaa ataacagaaa actaggacgt caaccaatta 2400
atatcgatgg ttcgaaagat acaactgatg gtaaaggctc tgaattctta ttagctaacg 2460
atattgacaa ctcaaactcg attgttcaag ctgagcaatt aaactggcta cactatttaa 2520
tgaatttttg tagtattaca ggtaataatg acaatgcgaa ttttgatggc attcgtgtag 2580
atgctgttga taatgttgat gctgatttac taaaaatagc tggcgattat tttaaagctc 2640
tatatggtac agataaaaagc gacgccaatg ccaataagca tttgtctatt ttagaagact 2700
ggaacggtaa agatcctcag tatgttaate aacagggcaa tgcgcaatta acaatggatt 2760
acacagttac ttcacagttt ggcaattctc taacacatgg cgccaacaac aggagtaaca 2820
tgtggatttt cttagatact ggctattatc ttaattggaga tcttaataag aagatagtag 2880
ataagaaccg tccaattctt tttgttagag cacatgatta cgatgctcaa gatccaatta 3000
aagttattcc aaattatagt tttgttagag cacatgatta cgatgctcaa gatccaatta 3000
gaaaagccat gattgatcat ggtattatta aaaacatgca ggatactttc acttttgacc 3060
aactggctca gggaatggaa ttctactata aagatcaaga gaatccgtct ggtttcaaaa 3120
agtataacga ttataactta cctagtgtt atgcaatgtt gttgactaat aaggatactg 3180
tacctcgtgt ctattatgga gatatgtacc tcaaggcgg gcaatatatg gaaaaagga 3240
cgatttacaa tctgtcatt tcagcgttgc tcaagctag aataaaatat gtttctggtg 3300
ggcaacaat ggctaccgat agttctggaa aagaccttaa agatggcgaa actgatttgt 3360
taacaagtgt tcgatttggt aaaggaatta tgacatcaga tcaaaccaca acacaagaca 3420
atagccaaga ttataaaaat caaggcatcg gtgtcattgt tggtaataac cctgacctta 3480
agttgaacaa tgataagacc attaccttgc atatgggaaa ggcgcataag aatcaacttt 3540
accgtgcctt agtattatca aatgactcag gaattgatgt ttatgatagt gatgataaag 3600
caccaacttt gagaacaaat gacaacggtg acttgatttt ccataagaca aatacgtttg 3660
tgaagcaaga tggaaactatt ataaattacg aaatgaaggg atcattaaat gcttttaatt 3720
caggttattt aggtgtctgg gtgccagttg gagctagtga ttcacaagat gctcgtacag 3780
tggcaactga gtcatcatca agtaatgat gttctgtatt ccattcaaat gctgcattag 3840
attctaattg tatatatgaa ggcttttcaa actttcaagc gatgccgact tctcctgagc 3900
aaagtacaaa tgttgattat gcaacaaagg ctaacttatt taaagaatta ggtattacta 3960
gttttgagtt agcacctcaa tataggtcta gtggtgacac taattacggg ggcatgtcat 4020
tcttagattc tttcttaaat aatggttatg catttaccga tagatatgat ttaggcttta 4080
caaaagcaga cgggaatcct aacccaacaa agtatggaac agatcaagat ttacgtaatg 4140
caatagaggc attacacaaa aacggcatgc aggctatagc tgattgggtt cctgacccaa 4200
tatatgcttt accagaaaag gaagttgta ccgctactag agtagacgaa cggggaaatc 4260
aactaaaaga cacagatttt gtcaacttac tctatgttgc taactactaa agtagtgggt 4320
tggattatca ggcaaagtat ggccggcgaat ttttagataa attaaagaaa gagtaccat 4380
cgttattcaa acagaaccaa gtatcgacag gtcagccaat tgatgcttct acaaaaaatta 4440

```

```

agcaatggtc agctaaatat atgaatggga ccaatatattt acatcgaggt gcttattatg 4500
ttttgaaaga ctgggctact aaccagtatt ttaacattgc aaaaacgaat gaagtatttt 4560
tgccactaca gttgcagaat aaagatgcgc aaactggttt cattagtgat gcctccgggtg 4620
taaaatatta ctcaattagt ggttatcaag caaaagatac ttttattgaa gatggtaatg 4680
ggaattggta ttactttgat aaagatgggt acatggtgcg ttcgcagcaa ggagaaaatc 4740
ctataagaac agtcgaaact agtgtcaaca cacgaaacgg taattattac tttatgccaa 4800
atgggtgtcga gttgcgcaaa ggctttggaa cggataatag tggtaatgtc tattattttg 4860
atgatcaagg taagatgggt agagataaat acattaacga tgatgctaata aattttttatc 4920
acttaaatgt tgatgggact atgtctcgag gactatttaa atttgattct gatactctac 4980
agtattttgc tagtaatgggt gtccaaataa aagatagtta tgcgaaggat agtaaaggca 5040
ataaatatta ttttgactca gctacaggaa ataacgatac tgggaaaagcc caaacttggg 5100
atggtaatgg ctactatatt actattgatt ctgatgcgaa caatacaatt ggggttaaca 5160
cagactacac tgcctacatc actagctcgc tgcgcgaaga tggcttattt gctaacgcac 5220
cttacgggtg tgtaacaaaa gaccaaaatg gtaacgatct taagtggcag tatattaacc 5280
atacgaaaca gtacgaaggg caacaagtgc aagtcacgcg tcaatacaca gacagtaagg 5340
gagtcagctg gaacttaatt acctttgctg gtggtgattt acaaggacaa aggctttggg 5400
tggtatgctg tgcgttaact atgacaccat ttaaaacgat gaaccaaata agcttcatta 5460
gttatgctaa ccgcaatgat gggttgtttt tgaatgcgcc ataccaagtc aaggggtatc 5520
aattagctgg gatgtccaac caatacaagg gccaaacaagt gaccattgct ggggtggcga 5580
acgtttctgg aaaagactgg agtctgatta gttttaatgg gacacagtac tggattgata 5640
gtcaggcatt gaataccaat ttcacacatg acatgaacca aaaggctctt gtcaatacaa 5700
ctagtaatct tgatgggtta ttcttaaatg cgccataccg tcaaccgggt tataagttag 5760
ccggtttggc taaaaattac aacaaccaaa cggttactgt tagtcaacag tactttgatg 5820
atcaaggcac ggtctggagt cagggtgtcc ttgggggtca gacggctctg gttgataacc 5880
atgcattggc acagatgcaa gttagtata cagaccaaca gctctatgtg aatagcaatg 5940
gtcggaaatga tgggttattc ttgaatgcgc catatcgtgg tcaagggtca caactgatag 6000
gcatgacggc agattataat gggcaacatg tacaagtac caagcaaggg caagatgcct 6060
atgggtgcaca atggcgctct attacgctaa ataatacaaa ggtctgggtt gatagtcgcg 6120
ctttgagcac aacaatcatg caagccatga atgataatat gtatgtaaat agcagccaac 6180
ggacagatgg cttgtgggta aacgcacctt atacgatgag tggggctaaa tgggctggtg 6240
atacacgttc agctaattgg cgctatgtcc atatttcaaa agcttattca aacgaagtcg 6300
gcaatacata ttacttgacg aatttgaatg gtcaaagcac atggattgac aagcgggctg 6360
ttactgtgac cttcgatcag gtgggtggcat taaatgcaac gattgtggca cgccaacgac 6420
cagatgggat gtttaagaca gcaccatat gtgaagcggg ggcgcagttt gtcgattatg 6480
tgacaaacta taaccagcaa accgtgccag taacaaagca acattcagat gctcagggga 6540
atcaatggta cttagcgaca gtgaatggga cacaatactg gattgatcaa cggtcatttt 6600
caccagtagt aacgaagggt gttgattatc aagctaagat tgtgccacgg acaacacgtg 6660
atgggtgtgt tagtggcgca ccctatgggg aagtgaatgc taagctagtt aacatggcaa 6720
ctgcgtatca aaatcaagtt gtccatgcga caggggaata tacgaatgct tcagggatca 6780
catggagtca gttcgcgtta agcgggcaag aagacaagct atggattgat aagcgtgctt 6840
tgcaagctta aggaaggat tcgacaaagg agggtaacat tatcagcgga tgggtgttatc 6900
ctcctttcct gtactcagta tttcccaaat aattgagaca gtttcatgac aaatcaacaa 6960
aactagtgtc aatgcctcgg ttatggggta aactactatt agttaaaggg ttgttgcata 7020
ataatatcac attcaatata ttatgtattt ttatctgatt atgtgatttt ttgagatttg 7080
gagcgaaaat gaataaagta ataattaata aaagtttctg tgttttggtg atgagtttgt 7140
tgtcgatatt tctattctca ttaagagtag acgctcgttc caatagtggc tacaatcaaa 7200
tatcttttaa tcaacatgaa attgcttatg cgctttttag tcaagtacct tggtatgcaa 7260
ctgtgaatgt cggaatgtct gcagacaaaa ataatatcta tacagctatt gatatggggc 7320
catagtatca aggaacttct tttttccgt ggtggtatca acatgataat tatgattatc 7380
atgatgg

```

&lt;210&gt; 18

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: artificial sequence

&lt;400&gt; 18

gggcccgccta gcatgaaaca acaagaaaca gt

32

&lt;210&gt; 19

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: artificial  
sequence

&lt;400&gt; 19

cccgggggtcg acctttgtcg aatccttccc

30

&lt;210&gt; 20

&lt;211&gt; 6204

&lt;212&gt; DNA

&lt;213&gt; Leuconostoc mesenteroides

&lt;400&gt; 20

```

gctagcatga aacaacaaga aacagttacc cgtaaaaaac tttataaatc cggtaagggtt 60
tgggttgacag cagctactgc atttgcggtg ttggggggtt caactgtaac aacagtccat 120
gcggtatacaa attcgaatgt cgctgttaag caaataaata atacaggaac caatgattct 180
ggcgaaaaaaa aggtaccggt tccatcaact aataatgata gtttgaagca aggaacagat 240
ggttttttggt atgattcaga cggcaatcgt gtcgatcaga agaccaatca gattctgctt 300
actgcggaac aacttaaaaa aaataacgaa aaaaatttat cagtaatcag tgatgatata 360
tcaaaaaaag atgatgaaaa tatttctaag cagacccaaa ttgctaatac acaaacagta 420
gatactgcta aaggcctgac taccagtaat ttatctgatc ccatcactgg gggtcactat 480
gaaaatcaca atggctactt tgtttatata gatgcttcag gaaaacaagt aacagggtttg 540
caaaatattg atggtaattt acaatatttt gatgacaatg gatatcaagt caagggatcc 600
ttccgagatg tcaacggcaa gcatatctat tttgattcag taacagggaa agctagtcca 660
aatgttgata ttgttaacgg taaagctcaa ggatatgatg cgcaaggcaa ccaattaaag 720
aaaagttatg tgcgccgatg ttctgggcaa acttactatt ttgatggtaa tggccaaccg 780
ttaatcggct tgcaaaacaat tgatgggaac ctacaatatt ttaaccaaca aggggttcaa 840
ataaagggtg gtttccaaga tgtaacaat aaacgtattt attttgacc aaacacaggt 900
aatgccgttg ccaataactg aataattaac ggtaaattac aggggcgtga cgcaaatggg 960
aaccaggtaa agaatgcatt tagtaaaagat gttgcaggaa atacatttta ttttgacgca 1020
aacggtgtga tgtaaacagg gttgcaaact atttcaggaa agacatatta tcttgatgaa 1080
caaggacacc tgagaaaaaa ttacgcggga acattcaata atcagtttat gtacttcgat 1140
gctgatacag gtgcgggtaa aacagcgatt gaatatcaat ttgatcaagg attggtatca 1200
caaagtaatg aaaatactcc tcacaatgcc gcaaagtctt atgataaaaag tagttttgaa 1260
aatgttgatg gttacttaac agcagataca tggatcgtc caaccgatat tttaaaaaat 1320
ggagatactt ggacggcatc taccgaaact gatatgcgtc cgcttttaat gacatggttg 1380
cctgacaaac aaacacaagc aaattacttg aattttatgt ctagttaaagg acttgggtata 1440
acgaccactt atacagcagc tacgtcacia aaaaacttaa atgacgcagc ctttggtatt 1500
caaacagcaa ttgaacaaca aatatctttg aaaaaaagta ctgagtggtt acgtgatgca 1560
attgatagtt ttgtgaagac gcaagctaatt tggaataagc aaacagaaga tgaagctttc 1620
gatggtttgc agtggcttca agggggattc ctagcttatc aagatgattc acatcggacg 1680
ccgaatactg attcaggaaa taacagaaaa ctaggacgtc aaccaattaa tatcgatggg 1740
tcgaaagata caactgatgg taaaggctct gaattcttat tagctaacga tattgacaac 1800
tcaaatccga ttgttcaagc tgagcaatta aactggctac actattttaat gaattttggt 1860
agtattacag gtaataatga caatgcgaat tttgatggca ttcgtgtaga tgctgttgat 1920
aatgttgatg ctgatttact aaaaatagct ggcgattatt ttaaagctct atatggtaca 1980
gataaaaagcg acgccaatgc caataagcat ttgtctattt tagaagactg gaacgggtaaa 2040
gatcctcagt atgttaatca acagggcaat gcgcaattaa caatggatta cacagttact 2100
tcacagtttg gcaattctct aacacatggc gccacaaca ggagtaacat gtgggtatttc 2160

```

ttagatactg	gctattatct	taatggagat	cttaataaga	agatagtaga	taagaaccgt	2220
ccaaattctg	gcactttggt	taacagaatt	gctaattcag	gtgatacaaa	agttattcca	2280
aattatagtt	ttgtagagc	acatgattac	gatgctcaag	atccaattag	aaaagccatg	2340
attgatcatg	gtattattaa	aaacatgcag	gatactttca	cttttgacca	actggctcag	2400
ggaatggaat	tctactataa	agatcaagag	aatccgtctg	gtttcaaaaa	gtataacgat	2460
tataaacttac	ctagtgccta	tgcaatggtg	ttgactaata	aggatactgt	acctcgtgtc	2520
tattatggag	atatgtacct	cgaaggcggg	caatatatgg	aaaaagggac	gatttacaat	2580
cctgtcattt	cagcgttgct	caaagctaga	ataaaatatg	tttctgggtg	gcaaacaatg	2640
gctaccgata	gttctggaaa	agaccttaaa	gatggcgaaa	ctgatttggt	aacaagtgtt	2700
cgatttggtg	aaggaattat	gacatcagat	caaaccacaa	cacaagacaa	tagccaagat	2760
tataaaaaatc	aaggcatcgg	tgctcattggt	ggtaataaacc	ctgaccttaa	gttgacaat	2820
gataagacca	ttaccttgca	tatgggaaaag	gcgcataaga	atcaacttta	ccgtgcctta	2880
gtattatcaa	atgactcagg	aattgatggt	tatgtagtg	atgataaagc	accaactttg	2940
agaacaaatg	acaacgggtg	cttgattttc	cataagacaa	atacgtttgt	gaagcaagat	3000
ggaactatta	taaattacga	aatgaaggga	tcattaaatg	ctttaatttc	aggttattta	3060
gggtgtctggg	tgccagttgg	agctagtgat	tcacaagatg	ctcgtacagt	ggcaactgag	3120
tcatcatcaa	gtaatgatgg	ttctgtattc	cattcaaag	ctgcattaga	ttctaattgt	3180
atatatgaag	gcttttcaaa	ctttcaagcg	atgccgactt	ctcctgagca	aagtacaaat	3240
gttggttattg	caacaaaggc	taacttattt	aaagaattag	gtattactag	ttttgagtta	3300
gcacctcaat	ataggtctag	tggtgacact	aattacgggtg	gcatgtcatt	cttagattct	3360
ttcttaaata	atggttatgc	atttaccgat	agatatgatt	taggcttta	caaagcagac	3420
gggaatccta	acccaacaaa	gtatggaaca	gatcaagatt	tacgtaatgc	aatagaggca	3480
ttacacaaaa	acggcatgca	ggctatagct	gattgggttc	ctgaccaa	atatgcttta	3540
ccaggaaagg	aagttgttac	cgctactaga	gtagacgaac	ggggaaatca	actaaaagac	3600
acagattttg	tcaacttact	ctatgttgct	aataactaaa	gtagtgggtg	ggattatcag	3660
gcaaagtatg	gcggcgcaat	tttagataaa	ttaagagaag	agtacccatc	gttattcaaa	3720
cagaaccaag	tatcgacagg	tcagccaatt	gatgcttcta	caaaaattaa	gcaatggtca	3780
gctaaatata	tgaatgggac	caatatttta	catcgagggtg	cttattatgt	tttgaaagac	3840
tgggctacta	accagtattt	taacattgca	aaaacgaatg	aagtattttt	gccactacag	3900
ttgcagaata	aagatgcgca	aactggtttc	attagtgatg	cctccgggtg	aaaatattac	3960
tcaattagtg	gttatcaagc	aaaagatact	tttattgaag	atggtaatgg	gaattgggtat	4020
tactttgata	aagatgggtta	catgggtcgt	tcgcagcaag	gagaaaatcc	tataagaaca	4080
gtcgaaacta	gtgtcaacac	acgaaacggg	aattattact	ttatgccaaa	tggtgtcgag	4140
ttgcgcaaa	gctttggaac	ggataatagt	ggtaattgtc	attattttga	tgatcaaggt	4200
aagatggtga	gagataaata	cattaacgat	gatgctaata	atttttatca	cttaaatgtt	4260
gatgggacta	tgctctgagg	actattttaa	tttgattctg	atactctaca	gtattttgct	4320
agtaatggtg	tcctaaataa	agatagtatt	gcgaaggata	gtaaaggcaa	taaataattat	4380
tttgactcag	ctacaggaaa	taacgatact	gggaaagccc	aaacttgga	tggtaatggc	4440
tactatatta	ctattgattc	tgatgcgaac	aatacaattg	gggttaacac	agactacact	4500
gcctacatca	ctagctcgct	gcgcgaagat	ggcttatttg	ctaacgcacc	ttacgggtgt	4560
gtaacaaaag	acaaaatgg	taacgatctt	aagtggcagt	atattaacca	tacgaaacag	4620
tacgaagggc	aacaagtgca	agtcacgcgt	caatacacag	acagtaaggg	agtcagctgg	4680
aacttaatta	cctttgctgg	tggtgattta	caaggacaaa	ggctttgggt	ggatagtcgt	4740
gcgttaacta	tgacaccatt	taaaacgatg	aaccaaataa	gcttcattag	ttatgctaac	4800
cgcaatgatg	ggttgttttt	gaatgcgcca	taccaagtca	aggggtatca	attagctggg	4860
atgtccaacc	aatacaaggg	ccaacaagtg	accattgctg	gggtggcgaa	cgtttctgga	4920
aaagactgga	gtctgattag	ttttaatggg	acacagtact	ggattgatag	tcaggcattg	4980
aataccaatt	tcacacatga	catgaaccaa	aaggtctttg	tcaatacaac	tagtaatctt	5040
gatgggttat	tcttaaatgc	gccataccgt	caaccgggtt	ataagttagc	cggtttggct	5100
aaaaattaca	acaaccaaac	ggttactgtt	agtcaacagt	actttgatga	tcaaggcacg	5160
gtctggagtc	aggttgtcct	tgggggtcag	acgggtctgg	ttgataacca	tgcatgggca	5220
cagatgcaag	ttagtgtac	agaccaacag	ctctatgtga	atagcaatgg	tcggaatgat	5280
gggttattct	tgaatgcgcc	atatcgtggg	caagggctac	aactgatagg	catgacggca	5340
gattataatg	ggcaacatgt	acaagtgacc	aagcaagggc	aagatgccta	tggtgcacaa	5400
tggtgtctta	ttacgctaaa	taatcaacag	gtctgggttg	atagtcgcgc	tttgagcaca	5460
acaatcatgc	aagccatgaa	tgataaatatg	tatgtaaat	gcagccaacg	gacagtggc	5520
ttgtgggttaa	acgcacctta	tacgatgagt	ggggtctaat	gggctggtga	tacacgttca	5580
gctaattgggc	gctatgtcca	tatttcaaaa	gcttattcaa	acgaagtcgg	caatacatat	5640
tacttgacga	atttgaatgg	tcaaagcaca	tggattgaca	agcgggctgt	tactgtgacc	5700
ttcgatcagg	tggtggcatt	aaatgcaacg	attgtggcac	gccaacgacc	agatgggatg	5760



```
tttaagacag caccatatgg tgaagcgggg gcgcagtttg tcgattatgt gacaaactat 5820
aaccagcaaa ccgtgccagt aacaaagcaa cattcagatg ctcaggggaa tcaatggtac 5880
ttagcgacag tgaatgggac acaatactgg attgatcaac ggtcattttc accagtagta 5940
acgaaggtgg ttgattatca agctaagatt gtgccacgga caacacgtga tgggtgtgtt 6000
agtggcgcac cctatgggga agtgaatgct aagctagtta acatggcaac tgcgtatcaa 6060
aatcaagttg tccatgcgac aggggaatat acgaatgctt cagggatcac atggagtcag 6120
ttcgcgtaaa gcgggcaaga agacaagcta tggattgata agcgtgcttt gcaagcttaa 6180
gggaaggatt cgacaaaggt cgac                                     6204
```

<210> 21

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 21

```
Asp Thr Asn Ser Asn Val Ala
  1                               5
```

<210> 22

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 22

```
Met Lys Gln Gln Glu
  1                               5
```

<210> 23

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 23

```
Lys Lys Val Pro Val
  1                               5
```

<210> 24

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 24

Lys Asp Asp Glu Asn  
1 5

<210> 25

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 25

Ile Asp Gly Asn Leu  
1 5

<210> 26

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 26

Tyr Val Ala Asp Ser  
1 5

<210> 27

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 27

His Leu Arg Lys Asn  
1 5

<210> 28

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 28

Asn Glu Asn Thr Pro  
1 5

<210> 29  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 29  
Asn Val Asp Gly Tyr  
1 5

<210> 30  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 30  
Asn Pro Asp Leu Lys  
1 5

<210> 31  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 31  
Ser Asn Asp Ser Gly  
1 5

<210> 32  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 32  
Asn Thr Phe Val Lys  
1 5

<210> 33  
<211> 5

<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 33  
Ile Ser Gly Tyr Leu  
1 5

<210> 34  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 34  
Ser Asn Ala Ala Leu  
1 5

<210> 35  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 35  
Arg Gln Tyr Thr Asp  
1 5

<210> 36  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 36  
Gln Leu Tyr Arg Ala  
1 5

<210> 37  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>

33/36

<223> Description of Artificial Sequence: artificial  
sequence

<400> 37

Asp Asp Lys Ala Pro  
1 5

<210> 38

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 38

Thr Arg Gln Tyr Thr  
1 5

<210> 39

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 39

Ile Thr Phe Ala Gly  
1 5

<210> 40

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 40

Asn Gln Tyr Lys Gly  
1 5

<210> 41

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial  
sequence

<400> 41

Leu Phe Leu Asn Ala  
1 5

<210> 42  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 42  
Gln Val Ser Asp Thr  
1 5

<210> 43  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 43  
Leu Ile Thr Leu Asn  
1 5

<210> 44  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 44  
Gly Arg Tyr Val His  
1 5

<210> 45  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 45  
Thr Ala Pro Tyr Gly  
1 5

<210> 46  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 46  
Val Val Asp Tyr Gln  
1 5

<210> 47  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 47  
Leu Ser Gly Gln Glu  
1 5

<210> 48  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<400> 48  
tctagactgc aaaatggcaa ctacta 26

<210> 49  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 49  
gtcgacgggtt tcatttggag tagtta 26

<210> 50  
<211> 276  
<212> DNA  
<213> Solanum tuberosum

<400> 50  
atggcaacta ctaaattcttt ttttaatttta ttttttatga tattagcaac tactagttca 60  
acatgtgcta agttggaaga aatgggttact gttctaagta ttgatggagg tggaattaag 120  
ggaatcattc cagctatcat tctcgaattt cttgaaggac aacttcagga agtggacaat 180  
aataaagatg caagacttgc agattacttt gatgtaattg gaggaacaag tacaggaggt 240

ttattgactg ctatgataac tactccaaat gaaacc

276

<210> 51  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 51  
gcgtactcta gacgtactcc gccatgacca c

31

<210> 52  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 52  
gcgtacgtcg acggccctga tgggtcccat

30

<210> 53  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 53  
ggccgggtcg acgatacaaa ttcgaatgtc gctg

34

<210> 54  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial  
sequence

<400> 54  
ggccgggtgc aggttacct cctttgtcga atc

33



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 August 2000 (17.08.2000)

PCT

(10) International Publication Number  
**WO 00/47727 A3**

- (51) International Patent Classification<sup>7</sup>: C12N 9/10, 15/54, 15/63, 15/70, 15/82, 5/10, A01H 5/00, C12P 21/02, 19/04, 19/18
- (21) International Application Number: PCT/EP00/00954
- (22) International Filing Date: 7 February 2000 (07.02.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
199 05 069.4 8 February 1999 (08.02.1999) DE
- (71) Applicants (for all designated States except US): PLANT-TEC BIOTECHNOLOGIE GMBH FORSCHUNG & ENTWICKLUNG [DE/DE]; Hermannswerder 14, D-14473 Potsdam (DE). MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN e.V. [DE/DE]; Berlin (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): KOSSMANN, Jens [DE/DE]; Düppelstr. 29, D-14163 Berlin (DE). WELSH, Thomas [DE/DE]; Schönburgstrasse 20, D-12103 Berlin (DE). QUANZ, Martin [DE/DE]; Oppelner Strasse 34, D-10997 Berlin (DE). KNUTH, Karola [DE/DE]; Schöneberger Str. 6, D-12103 Berlin (DE).
- (74) Agent: VOSSIUS & PARTNER; Siebertstrasse 4, D-81765 München (DE).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— With international search report.
- (88) Date of publication of the international search report:  
7 December 2000
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID MOLECULES ENCODING ALTERNANSUCRASE

(57) Abstract: Nucleic acid molecules encoding an alternansucrase are provided. Moreover, vectors, host cells and plant cells transformed by the herein-described nucleic acid molecules and plants containing them are provided. Furthermore, methods are described for preparing transgenic plants which synthesize the carbohydrate alternan, because of the insertion of nucleic acid molecules en-

# INTERNATIONAL SEARCH REPORT

Intern 1al Application No

PCT/EP 00/00954

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/10 C12N15/54 C12N15/63 C12N15/70 C12N15/82  
C12N5/10 A01H5/00 C12P21/02 C12P19/04 C12P19/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EPO-Internal, WPI Data, PAJ, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 13389 A (DU PONT ;CAIMI PERRY GERARD (US); HERSHEY HOWARD PAUL (US); KERR P) 18 May 1995 (1995-05-18) page 15, line 12 -page 17, line 20 page 24, line 3 - line 10 claims 1,7-10,18,24 ---	1-26
A	WO 96 04365 A (US GOVERNMENT) 15 February 1996 (1996-02-15) abstract; claim 4 ---	1
A	COTE G L ET AL: "PRODUCTION, ISOLATION, AND IMMOBILIZATION OF ALTERNANSUCRASE" CELLS ABSTRACTS PAPERS AMERICAN CHEMICAL SOCIETY MEETING,XX,XX, vol. 207, no. 1, 13 March 1994 (1994-03-13), XP002066232 the whole document ---	17,20-26
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

12 September 2000

Date of mailing of the international search report

20/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk

Authorized officer

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/00954

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	RAEMAEKERS M H M ET AL: "PRODUCTION OF ALTERNANSUCRASE BY LEUCONOSTOC MESENTEROIDES NRRL B-1355 IN BATCH FERMENTATION WITH CONTROLLED PH AND DISSOLVED OXYGEN" JOURNAL OF CHEMICAL TECHNOLOGY AND BIOTECHNOLOGY. (INTERNATIONAL JOURNAL OF BIOTECHNICAL AND CHEMICAL PROCESSES),GB,ELSEVIER APPLIED SCIENCE PUBLISHERS. BARKING, vol. 69, no. 4, 1 August 1997 (1997-08-01), pages 470-478, XP000702350 ISSN: 0268-2575 abstract page 471, left-hand column -----	1,17,20, 26
P,X	ARGUELLO-MORALES ET AL.: "Sequence analysis of the gene encoding alternansucrase, a sucrose glucosyltransferase from Leuconostoc mesenteroides NRRL B-1355" FEMS MICROBIOLOGY LETTERS, vol. 182, 1 January 2000 (2000-01-01), pages 81-85, XP000937860 abstract -----	1-10

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 00/00954

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9513389 A	18-05-1995	AU 1208795 A	29-05-1995
		BR 9408286 A	26-08-1997
		CA 2176109 A	18-05-1995
		EP 0728213 A	28-08-1996
		HU 75075 A	28-04-1997
		JP 9505467 T	03-06-1997
		PL 314296 A	02-09-1996
		US 5908975 A	01-06-1999
		ZA 9408867 A	09-05-1996
WO 9604365 A	15-02-1996	AU 697819 B	15-10-1998
		AU 3235295 A	04-03-1996
		CA 2196596 A	15-02-1996
		EP 0804545 A	05-11-1997
		US 5702942 A	30-12-1997
		US 5789209 A	04-08-1998